# PUBLIC HEALTH REPORTS

VOL. 48 DECEMBER 29, 1933 NO. 52

## EXPERIMENTAL STUDIES ON ACUTE MERCURIAL POISONING

By SANFORD M. ROSENTHAL, *Senior Pharmacologist, National Institute of Health, United States Public Health Service*

Previous attempts, under experimental conditions, to combat acute mercurial poisoning have been largely of no avail. Considerable work with sodium thiosulphate has yielded negative results (Haskell (1); Young and Taylor (2)). Recently Hesse (3) was able to protect a certain percentage of rabbits, rats, mice, and guinea pigs against a fatal subcutaneous dose of mercuric chloride by the use of strontium thioacetate, but Haskell and Forbes (4) showed that in dogs no such antidotal effect could be demonstrated following oral or subcutaneous intoxication. This fact was confirmed by Hesse himself.

We have studied several compounds under various conditions and have obtained one which can be shown, if properly administered, to protect rats and dogs against lethal doses of mercuric chloride.

For such a drug to be of benefit following intravenous administration, it is necessary for it to be comparatively stable in the body, to be of low toxicity, and at the same time to be able to exist in the body in a state which will react with mercury to form compounds of diminished toxicity. Excretion in the urine is desirable, as this may bring about a concentration of the substance in the kidney cells.

Sodium thiosulphate ($Na_2S_2O_3$) has previously been studied in this respect and found inadequate (5). As will be shown here, it is not capable of protecting kidney tissue against bichloride or of forming in the blood stream insoluble compounds with mercury. Other substances which have been used, such as calcium sulphide and sodium hydrosulphite ($Na_2S_2O_4$), while forming insoluble sulphides with great ease in the test tube, are so unstable in the body that they are broken down almost immediately after injection.

20288°—34——1

THE ANTAGONISM OF MERCURY ACTION AS SHOWN UPON THE OXYGEN CONSUMPTION OF EXCISED RAT TISSUES

*Sodium thiosulphate.*—The oxygen consumption of tissues *in vitro* was determined with the Warburg micro-respiration apparatus in a manner previously described (6). Rat tissues were suspended in Locke's solution containing 0.03 percent sodium bicarbonate and 0.2 percent glucose. All experiments were run at 37.6° C. in an atmosphere of air.

The behavior of thiosulphate revealed that while no protection was afforded to kidney tissue against the action of mercury, with




CHART 1.—The ability of thiosulphate to counteract the effect of $HgCl_2$ on rat testes if added before the mercury, but not if added later. Oxygen consumption of 100 mg rat testes in Locke's solution at 37.6° C. Atmosphere=air. Curve 1, testes alone. Curve 2, testes+m/500 thiosulphate. Curve 3, testes+m/500 thiosulphate+m/5000 $HgCl_2$. Curve 4, testes+$HgCl_2$, thiosulphate added in 10 minutes. Curve 5, testes+m/5000 $HgCl_2$.

other tissues its toxic effect can, under certain conditions, be completely antagonized.

Upon the oxygen consumption of rat testes, thiosulphate, when added first in amounts 10 times the molar quantity of the mercury, afforded complete protection against the toxic action of mercury. If the mercury was added first, and the thiosulphate later, no protection was observed (chart 1).

Upon the oxygen consumption of minced rat liver, thiosulphate gave a high degree of protection when added to the tissue either before or 5 minutes after the mercury was added (chart 2).

With the rat kidney, however, no protection could be obtained against the action of mercury, whether the thiosulphate was added before or after the mercury, and even when 20 times the molar

concentration of thiosulphate was used (chart 2). The basis for this lack of protection of renal tissue by thiosulphate is unknown, but it is possible that therein may lie the explanation for the selective nephrotoxic action of mercury.

*Experiments on glutathione.*—Glutathione, in its reduced state, is a sulphur compound related to cysteine, and also occurrs as a physiological constituent of animal tissues. Voegtlin, Dyer, and




CHART 2.—The antagonism of $HgCl_2$ by thiosulphate on liver tissue, and lack of effect on kidney. Oxygen consumption of 75 mg rat kidney. Curve 1, kidney. Curve 2, kidney+m/250 thiosulphate. Curve 3, kidney+m/5000 $HgCl_2$. Curve 4, kidney+m/5000 $HgCl_2$, m/250 thiosulphate 15 minutes later. Curve 5, kidney+thiosulphate, $HgCl_2$ added 2 minutes later. Curve 6, 100 mg rat liver. Curve 7, liver+m/250 thiosulphate. Curve 8, liver+m/250 thiosulphate, m/2500 $HgCl_2$ added in 2 minutes. Curve 9, liver+$HgCl_2$, thiosulphate added 5 minutes later. Curve 10, liver+$HgCl_2$.

Leonard (7) showed that the toxic action of arsenic could be counteracted by this compound, and further work on its chemical and physiological properties has been carried out in this laboratory (8, 9).

Glutathione is superior to thiosulphate in the protection of rat tissues *in vitro* against mercury action. This was manifested in that protection could be demonstrated when the glutathione was added 10 minutes or longer after the addition of the mercury, and

[1] The crystalline reduced glutathione used in these experiments was prepared by Dr. J. M. Johnson of this laboratory.

further in that this protection also existed for renal tissues. When from 5 to 10 times the molar quantity of glutathione [1] was used, as of mercuric chloride, the protection was almost complete, whether the glutathione was added either before or shortly after the mercury (chart 3). In the interpretation of chart 3, our previous work must be recalled (6) in which it was shown that glutathione, in the presence of renal tissue, slowly underwent oxidation, so that this added oxygen consumption must be taken into account in the experiments on kidney tissue. With other tissues glutathione




CHART 3.—The ability of SH glutathione to protect kidney tissue against mercury. Curve 1, oxygen consumption of 60 mg of rat kidney+m/500 glutathione. Curve 2, kidney+glutathione, m/5000 $HgCl_2$ in 2 minutes. Curve 3, kidney+$HgCl_2$, glutathione added 10 minutes later. Curve 4, kidney alone. Curve 5, kidney+m/5000 $HgCl_2$. Oxygen requirements of the glutathione=27 $mm^3$ of $O_2$.

in similar concentrations is kept largely reduced. In experiments on rat testes, some protection could be demonstrated from glutathione added up to 50 minutes following the bichloride.

*Formaldehyde sulphoxylate* ($NaHSO_2 \cdot CH_2O \cdot 2H_2O$, *rongalite, formopone.*)—Sodium formaldehyde sulphoxylate is a product formed by the union of sodium hydrosulphite ($Na_2S_2O_4$) and formaldehyde. The ensuing product is a powerful reducing agent which, however, is considerably less toxic than either of its components, and at the same time is much more stable in the animal organism. We have

recrystallized formaldehyde sulphoxylate from the technical product according to the following method supplied us by Dr. A. E. Sherndal, of Metz & Co.:

Two hundred grams are dissolved in 90 cc of water by gentle heating to 70° C. If the solution is not alkaline to litmus, add some sodium carbonate. There should be a formaldehyde odor to the solution. Filter through a hot suction funnel and cool down the filtrate in ice water. When all of the crystals have




CHART 4.—The ability of formaldehyde sulphoxylate to afford protection to kidney tissue against a subsequent addition of $HgCl_2$. Curve 1, 50 mg rat kidney+m/250 sulphoxylate. Curve 2, kidney alone. Curve 3, kidney+sulphoxylate, m/5000 $HgCl_2$ added in 3 minutes. Curve 4, kidney+$HgCl_2$, sulphoxylate added in 15 minutes. Curve 5, kidney+m/5000 $HgCl_2$. Similar results were obtained with 100 mg of rat testes.

precipitated, filter them off on a suction funnel, wash once with a small amount of cold water, and press out as dry as possible.

We have sealed these crystals in ampoules with the exclusion of air and found them stable for a period of weeks, at least. While the copper sulphate titration is the standard method of assay, we have found simple iodine titration of the aqueous solution after the method of Elvove (14) sufficiently accurate to detect deterioration. One hundred milligrams of the moist crystals dissolved in water should consume from 23 to 26 cc of 0.1 normal iodine, dependent on the amount of moisture.[2]

[2] We are indebted to Metz & Co., Merck & Co., and Diarsenol Co., for a supply of the sulphoxylate.

The ability of sulphoxylate to alter the reducing power of tissues and of the gastro-intestinal tract will be made the subject of another communication. Some results pertinent to the present investigation will be described in a later section of this paper.

Upon the oxygen consumption of rat testes and kidney, sulphoxylate afforded protection against mercury bichloride if the sulphoxylate was added first. No appreciable protection was observed if the sulphoxylate was added from 15 to 45 minutes following the mercury. The molar concentrations employed (m/250) were 20 times those of mercury. In these concentrations the sulphoxylate alone did not alter the oxygen uptake of the tissues, as shown in the control vessels set up for such determinations (chart 4). It is likely that in these *in vitro* experiments the protection afforded the excised rat tissues is largely concerned with the formation of insoluble mercury compounds, although we were unable to detect precipitates in the presence of the suspensions of tissue.

### THE ABILITY OF THIOSULPHATE AND GLUTATHIONE TO PROTECT SUBCUTANEOUS TISSUES AGAINST THE ACTION OF MERCURY

Another example of the ability of thiosulphate to protect certain tissues was shown in the prevention of the local inflammatory action of mercury when injected subcutaneously. When two hundredth molar bichloride of mercury was injected under the skin of the shaved ear of an albino rabbit, marked inflammation and local ulceration occurred. When this concentration of bichloride in tenth molar thiosulphate solution was used, slight or no inflammation and no ulceration followed. Equal volumes of the solutions of twice the desired strength were mixed immediately before injection, and no precipitate could be observed. Three injections of 0.1 cc each were made into each ear of 2 rabbits, 1 ear being employed as a control for the mercury alone. The same experiment was performed on two dogs with the injection of 0.3 cc at each site. Practically complete protection was also present in these animals.

Protection of subcutaneous tissues could also be demonstrated when a mixture of two hundredth molar bichloride of mercury and tenth molar glutathione (neutralized) were injected subcutaneously into the ears of rabbits and dogs. No precipitate could be seen when the solutions were mixed shortly before their injection. The technique employed was the same as with thiosulphate, and the results were essentially similar.

Because of the fact that sulphoxylate, when added to mercuric chloride in even very dilute solutions, results immediately in a heavy precipitate, it was not feasible to demonstrate an antagonism by subcutaneous injection.

ANTIDOTAL ACTION FOLLOWING THE SYSTEMIC ADMINISTRATION OF MERCURIC CHLORIDE TO ANIMALS

*Sodium thiosulphate.*—The minimum lethal dose of mercuric chloride when injected intravenously within 2 minutes' time into albino rats (Buffalo strain) was found to be 0.28 cc of a m/400 solution per 100 g of body weight. With this amount, 13 of 15 rats died in from 1 to 7 days, with an average time of 3.3 days. With a dose of 0.2 cc of m/400 bichloride per 100 g, there were no deaths in 7 rats (table 1).

It was soon found that the weight of the rats was an important factor in bichloride toxicity, as the susceptibility of the animals increased with weight. The age factor as shown by MacNider (10) may be important here. As far as possible, rats below 150 g weight were employed in the following experiments:

Six rats, from 128 to 138 g, were injected intravenously with 0.4 cc of n/10 thiosulphate per 100 g (57 times the molar quantity of mercury) immediately before the injection of a lethal dose of bichloride. All of these animals died (table 1). On autopsy the kidneys presented a gross appearance typical of mercurial nephritis. These results are consistent with the absence of protection afforded excised kidney tissue in the studies dealing with oxygen consumption.

In view of the several investigations with negative results previously reported on the use of thiosulphate in mercurial intoxication in dogs (1, 2, 5), no further work with this compound was done upon them.

*Glutathione.*—Confirmatory of the results obtained on the oxygen uptake of rat tissues, glutathione proved an effective antidote for mercury intoxication in rats, even when injected subsequent to an intravenous injection of mercury.

TABLE 1.—*Intravenous toxicity of $HgCl_2$ to rats; the ability of glutathione injected intravenously to protect rats against a lethal dose of $HgCl_2$; the ability of sulphoxylate to protect rats if injected prior to the $HgCl_2$; the lack of effect of sodium thiosulphate*

| Number of rats | Average weight | $HgCl_2$ | Antidote | Effect | |
|---|---|---|---|---|---|
| | | | | Survived | Died |
| 7 | 163 | 0.20 cc n/400 per 100 g intravenously | None | 7 | 0 |
| 3 | 153 | 0.28 cc n/400 per 100 g intravenously | do | 0 | 3 |
| 5 | 138 | do | do | 0 | 5 |
| 7 | 111 | do | do | 2 | 5 |
| 6 | 131 | do | Thiosulphate, 0.4 cc n/10 per 100 g before Hg. | 0 | 6 |
| 5 | 137 | do | Glutathione, 0.4 cc n/10 per 100 g before Hg. | 3 | 2 |
| 7 | 118 | do | do | 7 | 0 |
| 5 | 119 | do | Glutathione, 0.4 cc n/10 per 100 g ½ hour after Hg. | 4 | 1 |
| 7 | 118 | do | do | 7 | 0 |
| 10 | 193 | 0.32 cc n/400 per 100 g intravenously | None | 0 | 10 |
| 10 | 178 | do | Sulphoxylate, 1 g per kilogram 25 minutes after Hg. | 0 | 10 |
| 7 | 135.5 | 0.30 cc n/400 per 100 g intravenously | None | 0 | 7 |
| 7 | 145 | do | Sulphoxylate, 1 g per kilogram before Hg. | 5 | 2 |

One group of 5 rats, weighing from 122 to 142 g, was injected intravenously with 0.4 cc of n/10 glutathione (freshly neutralized) per 100 g immediately before the injection of a lethal dose of bichloride. Three of these animals survived. In another group of 7 rats, weighing from 112 to 130 g, similarly treated, all survived (table 1).

Two groups of rats were injected intravenously with a similar quantity of glutathione in one half hour following the injection of the mercury. In one series of 5 rats weighing from 108 to 124 g, 4 survived. In another series of 7 rats, weighing from 112 to 130 g, all survived (table 1).

An attempt to establish the minimum lethal dose of bichloride of mercury by mouth to rats revealed that these animals could tolerate such large doses that it was considered too irregular a method to employ. Five groups of 5 rats each were used, and from 20 to 70 mg of bichloride per kilogram were introduced into the stomach through a catheter. Only 1 or 2 of each group succumbed to doses above 30 mg per kilogram.

Because of the difficulty of obtaining sufficient amounts of glutathione, experiments on dogs are incomplete.

The minimum lethal dose of bichloride when injected intravenously into dogs has been established as 4 mg per kilogram body weight (4). We have employed this dose in 8 dogs, with 7 of the 8 succumbing in an average of 4.4 days.

Two dogs were given an intravenous injection of 0.45 g of freshly neutralized reduced glutathione (100 times the molar concentration of bichloride) per kilogram just before the injection of 4 mg of bichloride per kilogram. The animals died 12 and 14 hours later (table 2). Conclusions drawn from these experiments must be modified by the fact that the sample of glutathione was not highly purified and by the obvious paucity of material. The experiments of Hesse, for instance, with strontium thioacetate, on rats, guinea pigs, and rabbits, indicate that an optimum dose existed, beyond which much less or no protection against mercury was afforded.

It may be that with a larger number of animals and with varying doses of glutathione, some protection can be shown; but with the evidence at hand it must be concluded that while glutathione affords an extraordinary degree of protection against mercuric poisoning in rats, it behaves similarly to strontium thioacetate in that it is ineffective in dogs. Glutathione may be rapidly broken down when injected into dogs, as Abderhalden (11) was unable to recover any from the urine of dogs following the subcutaneous injection of 1 g.

The above evidence is somewhat strengthened by the results with cysteine. Two dogs were injected with 0.23 g (100 times the molar quantity of mercury) of freshly neutralized cysteine hydrochloride per kilogram just before the intravenous injection of mercury,

and both animals died, one in 14 hours and the other in 3 days (table 2). Cystine, the oxidized state of cysteine, is known of itself to produce kidney lesions when injected into or fed to animals.

*Formaldehyde sulphoxylate.*—Some pharmacological characteristics of this compound will be later reported in greater detail. Some of the experiments bearing upon the present problem may be summarized as follows:

The toxicity is quite low. Intraperitoneal injections of 1 g (10 percent solution) per kilogram into rats daily for 3 to 4 weeks produced no visceral changes and no symptoms, except that in some cases there was less rapid gain in weight than in control animals. There was evidence of severe local pain for 2 or 3 minutes at the site of the intraperitoneal injection. The single intravenous injection of 1 g per kilogram of body weight into rats, guinea pigs, rabbits, and dogs (10 to 20 percent solution) if administered slowly (2 to 4 minutes) was attended by no symptoms and no after effects that we have observed.

When given by mouth in doses up to 1 g per kilogram to rats and rabbits, no symptoms were observed except slight diarrhea in rats, which cleared up by the following day.

The stability of sulphoxylate in the body is shown by the following experiments:

When 1 g of sulphoxylate per kilogram is injected intravenously into rats or rabbits, it can be demonstrated in the blood serum for at least 5 hours after the injection. Sulphoxylate can be detected in the serum in 2 ways: (1) To 2 drops of serum add an excess of ammonium sulphate crystals and then 1 drop of a dilute solution of sodium nitroprusside. A green color results. (2) The second method is based on the great ability of sulphoxylate to reduce mercuric salts to insoluble black mercurous compounds and metallic mercury. To approximately 0.5 cc of undiluted serum (in a small test tube) add 2 or 3 drops of a 0.2 percent aqueous solution of mercuric chloride. Normally no precipitate results; but after the above-stated dose of sulphoxylate a precipitate is formed which turns black and settles to the bottom of the tube. In rabbits this reaction was still positive 5 hours after the injection but negative the next day. A sample of serum allowed to stand in an open test tube at room temperature for several days still gave a strongly positive reaction.

Following the intravenous injection of 1 g of sulphoxylate per kilogram into rabbits, the nitroprusside test on the urine was strongly positive for at least 10 hours later, but negative the next day. Tests upon the feces and lower intestinal contents were negative.

When rats were fed through a catheter 1 g of sulphoxylate (10 percent solution) per kilogram, nitroprusside tests for sulphoxylate 1 hour after administration were positive throughout the gastrointestinal tract as far as the rectum. The feces of another rat were strongly

positive 3 hours after administration. Tests on the urine of rats and rabbits made up to 6 hours after the oral dosage were positive. In dogs with bichloride poisoning that were given by mouth 0.5 to 1.0 g sulphoxylate per kilogram, the liquid stools were strongly positive for sulphoxylate 1 hour later. While sulphoxylate is less stable in acid solution than in alkaline, evidence that only a small proportion would be destroyed by the gastric acidity was demonstrated in that a 1 percent solution made up in 0.1 normal hydrochloric acid and kept at 37.6° C. showed 92 percent (by iodine titration) still present after 1½ hours, 80 percent after 3 hours, 73 percent after 5½ hours, and 57 percent in 23 hours.

If solutions of mercuric chloride and sulphoxylate are mixed in a test tube, a precipitate forms which rapidly blackens with the formation of mercurous compounds. Since no black precipitate is obtained with lead acetate, this action is not due to the presence of sulphide ions. Upon standing, the reduction may be shown to proceed as far as the formation of metallic mercury (12). Precipitates can be detected in aqueous solutions of sulphoxylate of 1 part in 300,000 when a few drops of 1 percent bichloride are added. Likewise a precipitate can be observed when a few drops of 1 percent sulphoxylate are added to 1 to 80,000 bichloride.

In the test tube sodium thiosulphate does not form a precipitate with mercuric chloride except in fairly high concentrations, and in animals no such precipitating action could be demonstrated in the serum 15 minutes after the intravenous injection of 1 g of thiosulphate per kilogram.

Experimental studies on the antagonism of mercuric poisoning by sulphoxylate in rats also conformed with the results obtained upon excised rat tissues. Protection of rats from an *intravenous* injection of bichloride occurred only if the sulphoxylate had been administered previously.

A dose of bichloride slightly larger than the M.L.D. was used and the one group of available rats was heavier than those previously employed, so that the test was more severe for the sulphoxylate. Ten rats averaging 178 g in weight received intravenously 0.32 cc of n/400 bichloride per 100 g. Twenty-five minutes later they were injected with 1 g of sulphoxylate per kilogram. All animals died, on an average, in 3 days. Ten control rats of an average weight of 193 g died, on an average, in 2.3 days.

Seven rats averaging 145 g in weight received an injection of 1 g of sulphoxylate per kilogram just before the injection of 0.3 cc of n/400 mercuric chloride. Five of the seven rats survived. Of 7 control rats, with an average weight of 135.5 g, all died, on an average, in 4.4 days (table 2).

TABLE 2.—*The ability of sulphoxylate injected intravenously to protect dogs against a lethal intravenous injection of $HgCl_2$; the lack of effect of glutathione and cysteine*

| Weight | $HgCl_2$ | Antidote | Effect |
|---|---|---|---|
| *Kg* | | | |
| 13.0 | 4 mg per kg intravenously | None | Dead in 1 day. |
| 15.5 | | | Dead in 10 days. |
| 10.0 | | | Dead in 4 days. |
| 8.0 | | | Survived. |
| 10.5 | | | Dead in 4 days. |
| 8.0 | | | Do. |
| 8.0 | | | Dead in 5 days. |
| 14.0 | | | Dead in 3 days. |
| 7.0 | 4 mg per kg intravenously | Glutathione, 0.45 g per kg before Hg | Dead in 12 hours. |
| 7.7 | | | Dead in 14 hours. |
| 7.0 | 4 mg per kg intravenously | Cysteine, 0.23 g per kg before Hg | Do. |
| 13.6 | | | Dead in 3 days. |
| 10.0 | 4 mg per kg intravenously | Sulphoxylate, 0.7 g per kg just before Hg. | Survived. |
| 10.5 | | | Do. |
| 12.0 | | | Do. |
| 18.0 | | | Do. |
| 6.0 | | | Do. |

Experiments on dogs revealed that protection could be afforded from a lethal intravenous dose of bichloride if preceded by an injection of sulphoxylate. It was also possible to save a high percentage of dogs from a lethal oral dose of mercury if proper sulphoxylate therapy was instituted an hour or more after administration of the bichloride. All dogs were kept under observation for at least a week before use in the following experiments.

Five dogs were injected intravenously with 0.7 g of sulphoxylate (40 percent solution) per kilogram and shortly afterward were given intravenously 4 mg of bichloride per kilogram. All animals survived. Of 8 control animals receiving bichloride alone, 7 died (table 2).

Our observations were next extended to the treatment of dogs following the oral administration of bichloride. The fatal dose by mouth for dogs is stated by Hesse (13) to be 35 mg per kilogram of body weight. Haskell and Forbes (4) place it at 20 mg per kilogram in fasting dogs that have received morphine to prevent vomiting.

Eight dogs from whom food was withheld for 18 hours were given 20 mg of morphine sulphate per kilogram subcutaneously, to prevent vomiting, and in approximately one half hour 20 mg of bichloride of mercury (1 percent solution) per kilogram by stomach tube, washed in with twice the volume of water. To 4 of the dogs was given intravenously 0.5 g of sulphoxylate (20 percent solution) per kilogram 17 to 34 minutes after the mercury, and again at 4½ hours after. To the control dogs were given intravenously 3 cc of 0.8 percent sodium chloride per kilogram approximately 4 hours after the mercury. Three of the four control animals died. None of the treated dogs died (table 3).

TABLE 3.—*The protective action of sulphoxylate injected intravenously subsequent to an oral dose of 20 mg of $HgCl_2$ per kilogram to dogs*

| Weight | $HgCl_2$ | Antidote | Interval after $HgCl_2$ | Effect |
|---|---|---|---|---|
| *Kg* | | | | |
| 9.3 | 20 mg per kilogram by mouth. | 0.8% NaCl, 3 cc per kilogram intravenously. | 4¼ hr | Died in 3 days. |
| 12.5 | | | 4 hr | Survived. |
| 20.7 | | | 3¾ hr | Died in 30 hours. |
| 16.0 | | | 4¼ hr | Died in 14 days. |
| 8.0 | 20 mg per kilogram by mouth. | Formaldehyde, sulphoxylate 0.5 g per kilogram intravenously. | 30 min. and 4½ hr | Survived. |
| 11.6 | | | 20 min. and 4½ hr | Do. |
| 11.1 | | | 34 min. and 4½ hr | Do. |
| 9.0 | | | 17 min. and 4½ hr | Do. |

Because of the depression produced by the above dose of morphine, an attempt was made to reduce the dosage to 10 to 15 mg per kilogram, but in some dogs this proved insufficient to produce quiescence and a further dose was required. It was also found more satisfactory to wait an hour after the morphine was given before administering the mercury. The animals were carefully watched for vomiting following oral administration of bichloride.

Three dogs received 35 mg of mercury per kilogram by mouth, followed by an intravenous injection of 0.5 g sulphoxylate (20 percent solution) in 30, 32, and 75 minutes; one dog survived. Three control dogs received a similar volume of salt solution (2.5 cc per kilogram) approximately 30 minutes after the mercury, with no survivals (table 4).

TABLE 4.—*Less beneficial effect of sulphoxylate given intravenously following a larger oral dose (25 to 35 mg per kilogram) of $HgCl_2$ to dogs*

| Weight | $HgCl_2$ | Antidote | Interval after $HgCl_2$ | Effect |
|---|---|---|---|---|
| *Kg* | | | | |
| 12.5 | 35 mg per kilogram by mouth. | 0.8% NaCl, 2.5 cc per kilogram intravenously. | ½ hr | Died in 2 days. |
| 14.0 | | | | Died in 1 hour. |
| 24.0 | | | | Died in 3 days. |
| 12.0 | ....do.... | Sulphoxylate 0.5 g per kilogram intravenously. | 32 min. | Died in 4 days. |
| 12.0 | | | 30 min. | Died in 3 days. |
| 11.4 | | | 75 min. | Survived. |
| 7.0 | 25 mg per kilogram by mouth. | ....do.... | 70 min. and 5 hr | Died in 3 days. |
| 14.0 | | | 60 min. and 6 hr | Died in less than 20 hours. |
| 9.3 | | | 33 min. and 6 hr | Survived. |

Seven dogs received 25 mg of bichloride per kilogram by mouth in the usual manner. Three of these animals were treated with two intravenous injections each of 0.5 g of sulphoxylate (20 percent) per kilogram. Two of the three died (table 4). Of four control dogs injected with equivalent volumes of salt solution at approximately the same intervals after the mercury, all died (table 5).

Of those animals that died following the administration of sulphoxylate by the intravenous route alone, histological examination of the kidneys by Dr. J. G. Pasternack, of this laboratory, revealed a strik-

ing difference between them and the control animals treated with salt solution. While the control animals showed extensive degenerative changes typical of acute mercurial nephritis, the treated animals showed only vascular congestion, and in some cases cloudy swelling and focal round cell infiltration.

On the other hand, it was found in both control and treated animals that the mucous membrane of the stomach and upper portion of the small intestine showed extensive necrotic changes, in most cases presenting on gross examination a black surface of necrotic tissue.

In view of previous experiments showing that no appreciable amounts of sulphoxylate could be found in the alimentary canal following its intravenous injection, it was believed that the gastrointestinal damage produced locally by the bichloride might be an important factor in the death of those animals receiving only intravenous therapy.

Accordingly, 18 other dogs, after the usual preparation, were given 25 mg of bichloride per kg by mouth. Twelve of these animals received 0.5 gm sulphoxylate (10–20 percent solution) per kg intravenously, and at the same time 0.5 to 1.0 gm per kilo (5 percent solution) through stomach tube, from 1 to 1½ hours following the mercury. Nine of these 12 animals survived and seemed to escape the toxic effects of bichloride, except some gastrointestinal inflammation, with diarrhea, and in some cases bloody stools, for several days. This represents injury which occurred prior to the therapy. The stools shortly after the treatment take on a dark greyish-brown color, which is due to the presence of the reduced mercury, and give a positive test for sulphoxylate.

Of the three treated animals that succumbed, two died in so short a time that kidney damage could not be responsible for the death. One (19 kg) died in 11 hours and showed at autopsy one lung partly consolidated and filled with a bloody serous fluid, while the other lung was normal; aspiration into the lung was an important factor in this fatality. The second dog (6.6 kg) was found dead the following morning (within 18 hours). Autopsy revealed extensive necrosis of the gastric mucosa and inflammation throughout the intestinal mucosa, while the kidneys histologically showed only diffuse cloudy swelling and vascular engorgement. The third dog (20 kg) lived for 8 days but refused all food during this time. Bloody stools and elevation of the blood nonprotein nitrogen were present during this period. Autopsy, however, showed no significant renal lesions on gross and histological examination. Pregnancy of about 1 month's development was found to be present.

The 6 control animals received 0.8 percent salt solution by mouth and intravenously in equivalent volume to that of sulphoxylate at approximately the same time after the mercury; 5 of the 6 died within

4 days. Another group of 4 dogs received this dose of bichloride by mouth and intravenous salt solution as therapy, with no survivals (table 5).

TABLE 5.—*The protection of dogs against an oral dose of 25 mg of $HgCl_2$ per kg when sulphoxylate is administered both by mouth and intravenously up to 90 minutes after the $HgCl_2$*

| Weight | $HgCl_2$ | Antidote | Interval after $HgCl_2$ | Effect |
|---|---|---|---|---|
| *Kg* | | | | |
| 8.2 | 25 mg per kg by mouth. | 0.8% NaCl, 2.5 cc per kg intravenously. | 1 hr. 7 min. and 5½ hr | Died in 4 days. |
| 14.1 | | | 1 hr., 6 hr | Died in 2 days. |
| 8.2 | | | do | Died in 20 days. |
| 12.1 | | | 40 min., 6 hr | Died in 2 days. |
| 16.0 | do | 0.8% NaCl, 2.5 cc per kg intravenously and 5.0 cc per kg by mouth. | 1 hr. 7 min | Died in 4 days. |
| 12.5 | | | 1 hr. 5 min | Survived. |
| 20.0 | | | 1 hr. 3 min | Died in 3 days. |
| 5.7 | | | 1 hr. 5 min | Died in 2 days. |
| 7.0 | | | 1 hr. 10 min | Died in 1 day. |
| 9.0 | | | 1 hr. 17 min | Died in 2 days. |
| 6.6 | do | 20% sulphoxylate, 0.5 g per kg intravenously and 1.0 g per kg by mouth. | 1 hr. 20 min | Died during night. |
| 10.5 | | | 1 hr. 10 min | Survived. |
| 7.4 | | | 1 hr. 7 min | Do. |
| 6.7 | | | 1 hr | Do. |
| 18.0 | | | 40 min | Do. |
| 7.4 | do | 10% sulphoxylate, 0.5 g per kg intravenously and 1.0 g per kg by mouth. | 1 hr. 5 min | Do. |
| 9.5 | | | 1 hr. 10 min | Do. |
| 19.0 | | | 1 hr. 30 min | Do. |
| 20.0 | | | 1 hr | Died in 8 days. |
| 19.0 | | | 1 hr. 15 min | Died in 11 hours. |
| 7.6 | | | 1 hr | Survived. |
| 8.4 | | | 1 hr. 10 min | Do. |

Determinations of blood nonprotein nitrogen were made upon these animals to obtain evidence of functional damage to the kidneys. Results upon the 12 animals treated with sulphoxylate showed in the 9 survivors no elevation of the nonprotein nitrogen throughout the period of observation, which extended up to 8 weeks. Two died before determinations could be made, while the dog that died in 8 days showed marked elevation up to the time of death (chart 5).

Of the 6 control dogs whose nonprotein nitrogen was studied, 4 showed striking increases up to the time of death. The fifth dog died before a determination could be made, while the sixth, the survivor in this group, showed only a slight increase (chart 5). Vomiting, which was not detected, probably occurred in this animal as a basis for the slight toxic effects.

## DISCUSSION

The favorable results of sulphoxylate therapy in dogs following oral intoxication with mercuric chloride suggest the usefulness of such treatment in human cases. Intravenous injections have been found to afford protection from kidney damage, while oral administration, by the reaction with the unabsorbed mercury to form insoluble and less toxic compounds, can undoubtedly give some local protection and also inhibit further absorption of mercury. The evidence obtained from rats and rat tissues indicates that after the mercury has combined with the cell protoplasm, no benefit is obtained from the use

of sulphoxylate. Considerable time is no doubt required after an oral dose of mercury before the full damage to the kidneys occurs. The length of time beyond that established by our experiments after which sulphoxylate therapy will be of benefit, remains to be determined.

Our experiments have established the low toxicity of sulphoxylate to animals. While small amounts are injected intravenously into human beings as an impurity in neoarsphenamine (up to 25 percent (14)), the injection into human beings of doses comparable to those employed in our animals has not heretofore been attempted. Because of the fact that strong solutions are irritating when injected subcutaneously, it cannot be administered in this manner.

We have had occasion up to the time of this report to try this therapy in only one human case.[3] This case is reported here primarily




CHART 5.—The ability of sulphoxylate to protect dogs from kidney damage following $HgCl_2$. The non-protein nitrogen in the blood of 5 control dogs and of 10 dogs treated with sulphoxylate; all received a lethal dose of $HgCl_2$ (25 mg per kilo) by mouth. Interrupted lines, control dogs receiving 0.8 percent NaCl as treatment. Continuous lines, dogs treated with sulphoxylate.

to demonstrate that large doses of sulphoxylate can also be tolerated by human beings without ill effects.

A man (C. L.), age 26, weight 52 kg (114 pounds), took four ½-gram tablets of mercuric chloride with suicidal intent. Vomiting did not occur until approximately 1½ hours later, and treatment was instituted at the hospital approximately 2 hours after the bichloride had been taken. His stomach was washed with 5 percent sulphoxylate. The washings showed considerable quantities of the dark gray reduced mercury. Three hundred cc of 5 percent sulphoxylate were left in the stomach; 250 cc of 10 percent sulphoxylate were then administered intravenously, the injection being slowly given over a period of 40 minutes. Vomiting occurred toward the end of the

[3] Since this was written we have treated 5 additional acute cases. All recovered without renal or intestinal lesions.

injection, due either to the treatment or to the mercury. A sample of blood was taken 20 minutes after the injection. The serum showed a strongly positive nitroprusside test for sulphoxylate; and upon the addition of 0.2 percent mercuric chloride, a grayish black precipitate was immediately formed. Following the therapy the patient had no symptoms other than abdominal discomfort. During the next day bloody stools were passed. Recovery was uneventful, with no albumin appearing in the urine and no elevation of the blood nonprotein nitrogen.

While the question of most suitable dosage in human beings remains to be established, we would suggest in the average case the following technique: Oral therapy of 10 to 15 g by stomach tube as described above, followed by 10 g of sulphoxylate as a 5 or 10 percent solution given intravenously, allowing at least 15 minutes for the injection, and a repetition of an intravenous dose of 5–10 g in 3 hours. The solutions for intravenous injection should be freshly prepared, and the sulphoxylate for this purpose should be a purified and recrystallized product. Sodium formaldehyde sulphoxylate can be obtained from manufacturers of neoarsphenamine.

It is not possible at present to say whether subsequent therapy will be of benefit in delayed symptoms of acute poisoning, or whether it will be of benefit in cases of chronic mercurialism.

In view of the ability of dilute solutions of sulphoxylate to reduce rapidly mercuric chloride to insoluble and less toxic mercurous compounds and to metallic mercury, its beneficial action in acute intoxication may be explained by its stability in the body and by the fact that intravenous injections confer upon the body fluids the power strongly to reduce mercuric salts. Thus it can persist unchanged throughout the alimentary canal following oral doses; it can remain for several hours in the blood following intravenous injections, in amounts capable of rapidly reducing added mercuric chloride; it can appear unchanged in the urine in considerable quantities, indicating that a concentration of this substance occurs in the kidney cells.

The observation that sodium thiosulphate can antagonize the action of mercury upon some tissues only under certain conditions and that it can afford no protection to kidney tissue should serve to delineate more clearly its field of use in the treatment of mercurialism.

SUMMARY

The ability of certain compounds to influence the toxicity of mercuric chloride has been studied by measuring the oxygen consumption of excised rat tissues. Sodium thiosulphate protected the rat testes against the action of mercury if added before the mercury, but not at all if the mercury was added first. With the liver, protection was observed when the thiosulphate was added either before or just after

the mercury. With the kidney, no protection could be observed in any case, whether the thiosulphate was added before or after. Injected along with bichloride subcutaneously into rabbits' ears, thiosulphate prevented the local inflammatory reaction of bichloride. Injected intravenously into rats, thiosulphate did not protect against a subsequent lethal injection of mercury.

Glutathione was able to counteract the toxic action of bichloride on the oxygen consumption of rat tissues, including kidney, when added either before or up to one-half hour after the addition of mercury. Glutathione prevented the local inflammatory reaction of bichloride when injected subcutaneously along with it. Glutathione saved 10 of 12 rats when injected intravenously previous to a lethal dose of bichloride. Eleven of twelve rats were also saved when the glutathione was injected one-half hour after the bichloride. Glutathione did not protect two dogs when injected previous to a lethal intravenous dose of bichloride. Similar results were obtained upon two dogs with cysteine.

Formaldehyde sulphoxylate counteracted the effect of mercuric chloride on the oxygen consumption of rat tissues if added before the bichloride, but not if added subsequently. Likewise, on rats injected intravenously with fatal doses of bichloride, 5 of 7 were saved if sulphoxylate was injected previously, but none of 10 if injected 25 minutes following the mercury.

Upon dogs the following results with sulphoxylate were obtained:

Five dogs received an intravenous injection of sulphoxylate prior to the injection of a fatal dose of bichloride; all survived. Of 8 control animals, 7 died. Of 4 dogs receiving 20 mg of bichloride per kg by mouth, and 2 intravenous injections of sulphoxylate subsequently (after 17 and 34 minutes, and again after 4½ hours), all survived. Of 4 control animals, 1 survived. Six dogs received 25 to 35 mg of bichloride per kg by mouth and intravenous injections of sulphoxylate subsequently. Two of these animals survived, while none of six controls survived. Twelve dogs received 25 mg of bichloride per kg by mouth and both oral and intravenous doses of sulphoxylate 40 to 90 minutes later; 9 of the 12 survived. Of a total of 10 control dogs receiving this dose of bichloride and intravenous and oral salt solution as therapy, 9 died. Nonprotein nitrogen determinations made subsequent to the intoxication on the treated dogs showed no elevation in 9 of 10 animals, while marked rises occurred in the control animals.

One human case of bichloride poisoning received intravenously 0.5 gm of sulphoxylate per kilogram of body weight and approximately 15 grams by stomach tube with no ill effects. Recovery without renal damage occurred in this case.

20288°—34——2

REFERENCES

(1) Haskell, C. C., Henderson, W. C., and Hamilton, J. R.: Jour. Am. Med. Assoc., **85**, 1808 (1925).
(2) Young, A. G., and Taylor, F. H. L.: Jour. Pharm. & Exp. Therap., **42**, 185 (1931).
(3) Hesse, E.: Arch. f. Exp. Path. u. Pharm., **144**, 327 (1929).
(4) Haskell, C. C., and Forbes, J. C.: Jour. Pharm. & Exp. Therap., **35**, 147 (1929).
(5) Melville, K. I., and Bruger, M.: Jour. Pharm. & Exp. Therap., **37**, 1 (1929).
(6) Rosenthal, S. M., and Voegtlin, C.: Pub. Health Rep., **46**, 521 (1931).
(7) Voegtlin, C., Dyer, H. A., and Leonard, C. S.: Pub. Health Rep., **38**, 1882 (1923).
(8) Voegtlin, C., Rosenthal, S. M., and Johnson, J. M.: Pub. Health Rep., **46**, 339 (1931).
(9) Voegtlin, C., Johnson, J. M., and Rosenthal, S. M.: Jour. Biol. Chem., **93**, 435 (1931).
(10) MacNider, W. deB.: Jour. Exp. Med., **36**, 1 (1917).
(11) Abderhalden, E., Buadze, S., and Geidel, W.: Fermentforschung, **13**, 147 (1932).
(12) Jellinek, K.: Das Hydrosulfit. F. Euke, Stuttgart, Pub. (1911).
(13) Hesse, E.: Arch. f. Exp. Path. u. Pharm., **117**, 266 (1926).
(14) Elvove, E.: Pub. Health Rep., **40**, 1235 (1925).

## COURT DECISION RELATING TO PUBLIC HEALTH

*Order of local manager of health and charity prohibiting sale of unpasteurized milk and cream held invalid where ordinance permitted such sale.*—(Colorado Supreme Court; *City and County of Denver et al.* v. *Gibson et al.*, 24 P. (2d) 751; decided July 3, 1933.) Under the provisions of the milk ordinance of Denver it was lawful for licensed dairymen to sell raw milk and cream of a certain standard in Denver. Section 4 of this ordinance authorized the manager of health and charity to formulate such regulations, not inconsistent with the ordinance, as were necessary to procure a standard of milk required by the ordinance. Purporting to act pursuant to such section 4, the manager of health and charity issued an order to the effect that after a certain date it would be unlawful to sell unpasteurized milk or cream. Suit to enjoin the enforcement of this order was brought by persons licensed to sell their dairy products in Denver, and the lower court granted an injunction. The case was taken to the supreme court, which body, in affirming the judgment of the trial court, said that "The conclusion is inevitable that the manager of health and charity assumed a legislative function and promulgated an order in derogation of an existing ordinance." The appellate tribunal quoted with approval the following language used by the trial judge:

* * * The ordinance permits the sale of milk, both raw and pasteurized, and establishes certain specific scientific standards therefor. Can the manager, in

effect, repeal the ordinance by an attempt to "ordain", as he has done by the express terms of the order, that it shall be unlawful to sell milk and cream on and after February 1, 1930, unless the same has been pasteurized, which directly contradicts the express terms of the ordinance? * * * My conclusion in the present case is that * * * the manager of health and charity was and is without power to make and enforce the order mentioned.

---

## DEATHS DURING WEEK ENDED DECEMBER 9, 1933

[From the Weekly Health Index, issued by the Bureau of the Census, Department of Commerce]

| | Week ended Dec. 9, 1933 | Corresponding week 1932 |
|---|---|---|
| Data from 85 large cities of the United States: | | |
| Total deaths | 8,565 | 8,644 |
| Deaths per 1,000 population, annual basis | 12.0 | 12.3 |
| Deaths under 1 year of age | 620 | 607 |
| Deaths under 1 year of age per 1,000 estimated live births (81 cities) | 53 | 51 |
| Deaths per 1,000 population, annual basis, first 49 weeks of year | 10.9 | 11.1 |
| Data from industrial insurance companies: | | |
| Policies in force | 67,326,257 | 69,666,314 |
| Number of death claims | 13,845 | 13,381 |
| Death claims per 1,000 policies in force, annual rate | 10.7 | 10.0 |
| Death claims per 1,000 policies, first 49 weeks of year, annual rate | 9.8 | 9.5 |

# PREVALENCE OF DISEASE

*No health department, State or local, can effectively prevent or control disease without knowledge of when, where, and under what conditions cases are occurring*

## UNITED STATES

### CURRENT WEEKLY STATE REPORTS

These reports are preliminary, and the figures are subject to change when later returns are received by the State health officers

**Reports for Weeks Ended Dec. 16, 1933, and Dec. 17, 1932**

*Cases of certain communicable diseases reported by telegraph by State health officers for weeks ended Dec. 16, 1933, and Dec. 17, 1932*

| Division and State | Diphtheria | | Influenza | | Measles | | Meningococcus meningitis | |
|---|---|---|---|---|---|---|---|---|
| | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 |
| **New England States:** | | | | | | | | |
| Maine | 3 | 3 | 15 | ---- | 1 | ---- | 0 | 0 |
| New Hampshire | 1 | 1 | ---- | ---- | 12 | 1 | 0 | 0 |
| Vermont | 3 | 2 | ---- | ---- | 59 | ---- | 0 | 0 |
| Massachusetts [1] | 26 | 40 | ---- | 10 | 482 | 110 | 1 | 1 |
| Rhode Island | 2 | 4 | ---- | 1 | 9 | ---- | 1 | 0 |
| Connecticut | 10 | 5 | 4 | 13 | 17 | 13 | 0 | 0 |
| **Middle Atlantic States:** | | | | | | | | |
| New York | 54 | 49 | [2] 28 | [2] 45 | 584 | 715 | 3 | 3 |
| New Jersey | 25 | 41 | 20 | 32 | 99 | 230 | 0 | 0 |
| Pennsylvania | 51 | 92 | ---- | ---- | 327 | 208 | 3 | 2 |
| **East North Central States:** | | | | | | | | |
| Ohio | 65 | 78 | 101 | 644 | 120 | 203 | 3 | 3 |
| Indiana | 65 | 67 | 61 | 1,078 | 39 | 24 | 2 | 2 |
| Illinois | 52 | 80 | 11 | 167 | 34 | 54 | 4 | 11 |
| Michigan | 26 | 31 | 4 | 57 | 37 | 271 | 0 | 3 |
| Wisconsin | 13 | 13 | 17 | 111 | 161 | 222 | 3 | 0 |
| **West North Central States:** | | | | | | | | |
| Minnesota | 8 | 6 | ---- | 10 | 8 | 84 | 1 | 3 |
| Iowa [3] | 18 | 27 | 1 | ---- | 30 | 3 | 1 | 0 |
| Missouri | 80 | 26 | 6 | 184 | 112 | 14 | 2 | 4 |
| North Dakota | 10 | 9 | 2 | ---- | 33 | 120 | 1 | 0 |
| South Dakota | 22 | 20 | ---- | 17 | 217 | 2 | 0 | 0 |
| Nebraska | 6 | 35 | ---- | 26 | 13 | 1 | 1 | 1 |
| Kansas | 29 | 28 | 1 | 41 | 43 | 6 | 0 | 2 |
| **South Atlantic States:** | | | | | | | | |
| Delaware | 1 | 4 | ---- | 1 | 1 | 1 | 1 | 0 |
| Maryland [3] | 25 | 26 | 20 | 171 | 21 | 6 | 0 | 1 |
| District of Columbia | 10 | 5 | 1 | 64 | 25 | ---- | 1 | 2 |
| Virginia | 67 | 39 | ---- | ---- | 87 | 147 | 0 | 0 |
| West Virginia | 47 | 25 | 14 | 62 | 6 | 88 | 2 | 4 |
| North Carolina | 60 | 38 | 14 | 192 | 503 | 49 | 2 | 0 |
| South Carolina [1] | 14 | 12 | 459 | 1,446 | 125 | 11 | 0 | 0 |
| Georgia [1] | 35 | 36 | ---- | 3,954 | 299 | ---- | 1 | 0 |
| Florida | 22 | 23 | 3 | 38 | ---- | 1 | 1 | 1 |

See footnotes at end of table.

*Cases of certain communicable diseases reported by telegraph by State health officers for weeks ended Dec. 16, 1933, and Dec. 17, 1932*—Continued

| Division and State | Diphtheria | | Influenza | | Measles | | Meningococcus meningitis | |
|---|---|---|---|---|---|---|---|---|
| | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 |
| East South Central States: | | | | | | | | |
| Kentucky | 60 | 29 | 25 | 2,537 | 4 | | 1 | 1 |
| Tennessee | 44 | 25 | 113 | 3,767 | 269 | 5 | 1 | 2 |
| Alabama [1] | 33 | 27 | 47 | 7,034 | 114 | 3 | 1 | 2 |
| Mississippi [3] | 19 | 6 | | | | | 0 | 1 |
| West South Central States: | | | | | | | | |
| Arkansas | 18 | 13 | 32 | 4,272 | 294 | 1 | 0 | 0 |
| Louisiana [1] | 30 | 26 | 11 | 4,945 | 1 | | 0 | 2 |
| Oklahoma [4] | 70 | 29 | 53 | 2,305 | 39 | 2 | 0 | 0 |
| Texas [1] | 207 | 104 | 143 | 498 | 193 | 232 | 0 | 0 |
| Mountain States: | | | | | | | | |
| Montana | 7 | 1 | 5 | 1,388 | 2 | 449 | 2 | 1 |
| Idaho | | 5 | | 9 | 10 | 5 | 0 | 1 |
| Wyoming | | | | 101 | 34 | 17 | 0 | 0 |
| Colorado | 12 | 6 | | 313 | 4 | 8 | 0 | 0 |
| New Mexico | 10 | 10 | | 8 | 74 | 1 | 0 | 0 |
| Arizona | 5 | 5 | 20 | 174 | 3 | | 0 | 1 |
| Utah [3] | | 2 | 2 | 21 | 129 | | 1 | 1 |
| Pacific States: | | | | | | | | |
| Washington | 6 | 2 | 3 | 1 | 219 | 5 | 0 | 0 |
| Oregon | 1 | 1 | 17 | 769 | 18 | 45 | 0 | 1 |
| California | 32 | 64 | 48 | 1,271 | 137 | 27 | 3 | 1 |
| Total | 1,404 | 1,220 | 1,301 | 37,777 | 5,048 | 3,384 | 43 | 57 |

| Division and State | Poliomyelitis | | Scarlet fever | | Smallpox | | Typhoid fever | |
|---|---|---|---|---|---|---|---|---|
| | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 |
| New England States: | | | | | | | | |
| Maine | 0 | 0 | 10 | 31 | 0 | 0 | 1 | 0 |
| New Hampshire | 1 | 0 | 12 | 21 | 0 | 0 | 1 | 0 |
| Vermont | 0 | 0 | 15 | 11 | 0 | 0 | 3 | 0 |
| Massachusetts [1] | 0 | 2 | 222 | 360 | 0 | 0 | 3 | 7 |
| Rhode Island | 0 | 0 | 17 | 34 | 0 | 0 | 1 | 0 |
| Connecticut | 0 | 0 | 63 | 63 | 0 | 0 | 0 | 2 |
| Middle Atlantic States: | | | | | | | | |
| New York | 7 | 1 | 466 | 594 | 0 | 3 | 8 | 8 |
| New Jersey | 1 | 3 | 146 | 213 | 0 | 0 | 5 | 5 |
| Pennsylvania | 1 | 6 | 418 | 651 | 0 | 0 | 12 | 7 |
| East North Central States: | | | | | | | | |
| Ohio | 4 | 2 | 553 | 550 | 2 | 8 | 7 | 21 |
| Indiana | 1 | 0 | 183 | 118 | 4 | 5 | 4 | 4 |
| Illinois | 1 | 0 | 379 | 391 | 3 | 2 | 7 | 11 |
| Michigan | 1 | 0 | 293 | 297 | 0 | 0 | 11 | 6 |
| Wisconsin | 1 | 2 | 101 | 84 | 64 | 1 | 0 | 1 |
| West North Central States: | | | | | | | | |
| Minnesota | 0 | 0 | 100 | 73 | 3 | 0 | 2 | 0 |
| Iowa [3] | 0 | 0 | 87 | 47 | 1 | 64 | 4 | 0 |
| Missouri | 0 | 0 | 131 | 76 | 3 | 0 | 1 | 1 |
| North Dakota | 0 | 2 | 34 | 12 | 0 | 0 | 1 | 0 |
| South Dakota | 0 | 0 | 11 | 10 | 0 | 1 | 0 | 2 |
| Nebraska | 0 | 0 | 28 | 50 | 2 | 3 | 5 | 0 |
| Kansas | 0 | 1 | 115 | 88 | 7 | 1 | 5 | 0 |
| South Atlantic States: | | | | | | | | |
| Delaware | 1 | 0 | 6 | 11 | 0 | 0 | 1 | 0 |
| Maryland [3] | 0 | 0 | 80 | 100 | 0 | 0 | 9 | 7 |
| District of Columbia | 0 | 0 | 14 | 12 | 0 | 0 | 1 | 0 |
| Virginia | 0 | 0 | 128 | 78 | 0 | 0 | 16 | 13 |
| West Virginia | 0 | 0 | 144 | 69 | 0 | 0 | 7 | 5 |
| North Carolina | 0 | 1 | 131 | 77 | 1 | 0 | 6 | 4 |
| South Carolina [1] | 0 | 0 | 21 | 13 | 0 | 2 | 6 | 5 |
| Georgia [1] | 3 | 2 | 27 | 22 | 0 | 0 | 11 | 2 |
| Florida | 0 | 0 | 5 | 8 | 0 | 0 | 3 | 5 |

See footnote at end of table.

*Cases of certain communicable diseases reported by telegraph by State health officers for weeks ended Dec. 16, 1933, and Dec. 17, 1932*—Continued

| Division and State | Poliomyelitis | | Scarlet fever | | Smallpox | | Typhoid fever | |
|---|---|---|---|---|---|---|---|---|
| | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 | Week ended Dec. 16, 1933 | Week ended Dec. 17, 1932 |
| East South Central States: | | | | | | | | |
| Kentucky | 0 | 0 | 114 | 40 | 0 | 1 | 5 | 7 |
| Tennessee | 0 | 2 | 129 | 39 | 2 | 19 | 6 | 12 |
| Alabama [1] | 1 | 0 | 37 | 20 | 0 | 1 | 4 | 0 |
| Mississippi [3] | 1 | 0 | 25 | 12 | 3 | 1 | 0 | 1 |
| West South Central States: | | | | | | | | |
| Arkansas | 0 | 0 | 15 | 23 | 2 | 2 | 3 | 5 |
| Louisiana [1] | 1 | 0 | 14 | 12 | 30 | 0 | 15 | 14 |
| Oklahoma [4] | 1 | 0 | 47 | 34 | 0 | 4 | 5 | 0 |
| Texas [1] | 0 | 0 | 122 | 82 | 12 | 7 | 35 | 6 |
| Mountain States: | | | | | | | | |
| Montana | 0 | 0 | 15 | 10 | 18 | 0 | 2 | 1 |
| Idaho | 0 | 0 | 8 | 4 | 1 | 1 | 0 | 1 |
| Wyoming | 0 | 0 | 12 | 13 | 0 | 0 | 0 | 0 |
| Colorado | 0 | 0 | 19 | 25 | 3 | 0 | 8 | 0 |
| New Mexico | 0 | 0 | 33 | 18 | 0 | 0 | 6 | 1 |
| Arizona | 0 | 0 | 13 | 5 | 0 | 0 | 1 | 1 |
| Utah [3] | 0 | 0 | 10 | 25 | 11 | 0 | 0 | 0 |
| Pacific States: | | | | | | | | |
| Washington | 4 | 0 | 37 | 44 | 2 | 22 | 3 | 0 |
| Oregon | 1 | 0 | 44 | 20 | 9 | 7 | 2 | 2 |
| California | 6 | 3 | 205 | 111 | 8 | 1 | 20 | 8 |
| Total | 37 | 27 | 4,831 | 4,701 | 191 | 156 | 255 | 17 |

[1] Typhus fever, week ended Dec. 16, 1933, 54 cases, as follows: Massachusetts, 1; South Carolina, 15 Georgia, 22; Alabama, 12; Louisiana, 1, Texas, 17.
[2] New York City only.
[3] Week ended earlier than Saturday.
[4] Exclusive of Oklahoma City and Tulsa.

## SUMMARY OF MONTHLY REPORTS FROM STATES

The following summary of cases reported monthly by States is published weekly and covers only those States from which reports are received during the current week.

| State | Meningococcus meningitis | Diphtheria | Influenza | Malaria | Measles | Pellagra | Poliomyelitis | Scarlet fever | Smallpox | Typhoid fever |
|---|---|---|---|---|---|---|---|---|---|---|
| *October 1933* | | | | | | | | | | |
| Vermont | | 2 | | | 3 | | | 50 | 0 | 1 |
| *November 1933* | | | | | | | | | | |
| District of Columbia | 2 | 89 | 3 | | 49 | 1 | 0 | 58 | 0 | 17 |
| Indiana | 8 | 507 | 216 | | 93 | | 3 | 875 | 16 | 7 |
| Maine | 1 | 13 | 2 | | 9 | | 8 | 48 | 0 | 12 |
| Michigan | 6 | 116 | 7 | 4 | 170 | | 3 | 1,169 | 3 | 39 |
| Minnesota | 2 | 80 | 2 | | 127 | | 19 | 301 | 15 | 15 |
| Nebraska | | 39 | 27 | | 32 | | 5 | 175 | 24 | 5 |
| New Hampshire | | 3 | | | | | 2 | 85 | 0 | 0 |
| New Jersey | 5 | 117 | 71 | 1 | 130 | | 4 | 509 | 0 | 21 |
| New Mexico | 1 | 42 | 16 | 31 | 99 | 1 | 2 | 102 | 0 | 65 |
| North Carolina | 10 | 529 | 82 | | 806 | 22 | 6 | 840 | 1 | 24 |
| Ohio | 3 | 426 | 256 | 2 | 402 | | 16 | 2,287 | 4 | 44 |
| Tennessee | 4 | 310 | 186 | 391 | 574 | 7 | 6 | 539 | 9 | 75 |
| Texas | 1 | 1,231 | 578 | | 71 | 45 | 8 | 381 | 31 | 175 |
| Vermont | | 12 | | | 161 | | 3 | 52 | 0 | 0 |

| *October 1933* | Cases |
|---|---|
| Vermont: | |
| Chicken pox | 71 |
| Mumps | 23 |
| Undulant fever | 1 |
| Whooping cough | 83 |
| *November 1933* | |
| Anthrax: | |
| Texas | 1 |
| Chicken pox: | |
| District of Columbia | 43 |
| Indiana | 590 |
| Maine | 269 |
| Michigan | 1,564 |
| Minnesota | 1,124 |
| Nebraska | 297 |
| New Jersey | 910 |
| New Mexico | 28 |
| North Carolina | 295 |
| Ohio | 2,298 |
| Tennessee | 63 |
| Texas | 105 |
| Vermont | 278 |
| Conjunctivitis: | |
| New Mexico | 5 |
| Dengue: | |
| Texas | 6 |
| Diarrhea and enteritis: | |
| Ohio (under 2 years) | 17 |
| Dysentery: | |
| Michigan | 26 |
| Minnesota (amœbic) | 24 |
| Minnesota (bacillary) | 2 |
| New Jersey (amœbic) | 8 |
| New Mexico | 5 |
| Ohio | 19 |
| Tennessee | 5 |
| Texas | 45 |
| Food poisoning: | |
| New Mexico | 1 |
| Ohio | 13 |
| German measles: | |
| Maine | 8 |
| Michigan | 34 |
| New Jersey | 10 |
| New Mexico | 11 |
| North Carolina | 8 |
| Ohio | 17 |
| Tennessee | 2 |
| Impetigo contagiosa: | |
| Tennessee | 17 |
| Jaundice, spirochetal: | |
| Michigan | 8 |
| Lead poisoning: | |
| Ohio | 21 |
| Lethargic encephalitis: | |
| District of Columbia | 1 |
| Indiana | 3 |
| Maine | 1 |
| Michigan | 5 |
| Minnesota | 5 |
| Nebraska | 1 |
| New Jersey | 5 |
| New Mexico | 1 |
| Ohio | 8 |
| Tennessee | 5 |
| Texas | 9 |
| Mumps: | |
| Indiana | 24 |
| Maine | 17 |
| Michigan | 284 |
| Nebraska | 27 |
| New Jersey | 164 |
| New Mexico | 21 |
| Ohio | 108 |
| Tennessee | 41 |
| Texas | 25 |
| Vermont | 23 |
| Ophthalmia neonatorum: | |
| New Jersey | 3 |
| Ohio | 75 |
| Tennessee | 1 |
| Texas | 1 |
| Paratyphoid fever: | |
| Michigan | 3 |
| Minnesota | 2 |
| Ohio | 1 |
| Tennessee | 2 |
| Texas | 3 |
| Puerperal septicemia: | |
| New Mexico | 2 |
| Ohio | 5 |
| Tennessee | 2 |
| Rabies in animals: | |
| Indiana | 27 |
| New Jersey | 12 |
| Rabies in man: | |
| Ohio | 1 |
| Rocky Mountain spotted fever: | |
| North Carolina | 1 |
| Scabies: | |
| Tennessee | 4 |
| Septic sore throat: | |
| Maine | 1 |
| Michigan | 59 |
| Nebraska | 2 |
| North Carolina | 12 |
| Ohio | 211 |
| Tennessee | 18 |
| Vermont | 1 |
| Tetanus: | |
| Michigan | 1 |
| Minnesota | 1 |
| Ohio | 1 |
| Tennessee | 1 |
| Texas | 1 |
| Trachoma: | |
| New Jersey | 1 |
| Ohio | 2 |
| Tennessee | 13 |
| Texas | 6 |
| Trichinosis: | |
| New Jersey | 2 |
| Tularaemia: | |
| Michigan | 5 |
| Minnesota | 6 |
| Ohio | 7 |
| Texas | 1 |
| Typhus fever: | |
| North Carolina | 5 |
| Texas | 20 |
| Undulant fever: | |
| Indiana | 1 |
| Maine | 2 |
| Michigan | 4 |
| Minnesota | 12 |
| Nebraska | 1 |
| New Jersey | 1 |
| Ohio | 6 |
| Tennessee | 2 |
| Texas | 11 |
| Vincent's infection: | |
| Maine | 2 |
| Michigan | 24 |
| Tennessee | 2 |
| Whooping cough: | |
| District of Columbia | 56 |
| Indiana | 119 |
| Maine | 213 |
| Michigan | 834 |
| Minnesota | 233 |
| Nebraska | 279 |
| New Jersey | 485 |
| New Mexico | 96 |
| North Carolina | 530 |
| Ohio | 861 |
| Tennessee | 120 |
| Texas | 277 |
| Vermont | 210 |

## WEEKLY REPORTS FROM CITIES

*City reports for week ended Dec. 9, 1933*

| State and city | Diphtheria cases | Influenza cases | Influenza deaths | Measles cases | Pneumonia deaths | Scarlet fever cases | Smallpox cases | Tuberculosis deaths | Typhoid fever cases | Whooping cough cases | Deaths all causes |
|---|---|---|---|---|---|---|---|---|---|---|---|
| Maine: | | | | | | | | | | | |
| Portland | 0 | ------ | 0 | 0 | 10 | 5 | 0 | 1 | 0 | 3 | 27 |
| New Hampshire: | | | | | | | | | | | |
| Concord | 0 | ------ | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 6 |
| Manchester | 0 | ------ | 0 | 0 | 1 | 3 | 0 | 1 | 0 | 0 | 15 |
| Nashua | 0 | ------ | 0 | 0 | 0 | 3 | 0 | 0 | 0 | 0 | 0 |
| Vermont: | | | | | | | | | | | |
| Barre | 0 | ------ | 0 | 66 | 0 | 0 | 0 | 1 | 0 | 0 | 3 |
| Burlington | 1 | ------ | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 2 | 9 |
| Massachusetts: | | | | | | | | | | | |
| Boston | 7 | ------ | 0 | 146 | 20 | 46 | 0 | 6 | 0 | 50 | 217 |
| Fall River | 3 | ------ | 0 | 2 | 2 | 0 | 0 | 0 | 2 | 0 | 25 |
| Springfield | 0 | 1 | 1 | 0 | 3 | 1 | 0 | 0 | 0 | 16 | 36 |
| Worcester | 1 | ------ | 0 | 348 | 9 | 11 | 0 | 1 | 2 | 16 | 49 |
| Rhode Island: | | | | | | | | | | | |
| Pawtucket | 0 | ------ | 0 | 0 | 0 | 3 | 0 | 0 | 0 | 0 | 0 |
| Providence | 0 | ------ | 0 | 0 | 5 | 9 | 0 | 4 | 0 | 28 | 62 |
| Connecticut: | | | | | | | | | | | |
| Bridgeport | 0 | ------ | 0 | 5 | 2 | 7 | 0 | 1 | 0 | 2 | 34 |
| Hartford | 1 | ------ | 0 | 1 | 1 | 11 | 0 | 1 | 0 | 2 | 41 |
| New Haven | 0 | ------ | 0 | 0 | 1 | 4 | 0 | 0 | 0 | 2 | 45 |

*City reports for week ended Dec. 9, 1933*—Continued

| State and city | Diphtheria cases | Influenza | | Measles cases | Pneumonia deaths | Scarlet fever cases | Smallpox cases | Tuberculosis deaths | Typhoid fever cases | Whooping cough cases | Deaths all causes |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Cases | Deaths | | | | | | | | |
| New York: | | | | | | | | | | | |
| Buffalo | | | | | | | | | | | |
| New York | 37 | 22 | 13 | 29 | 163 | 130 | 0 | 83 | 6 | 101 | 1,542 |
| Rochester | 0 | | 1 | 0 | 9 | 9 | 0 | 0 | 0 | 7 | 73 |
| Syracuse | 0 | | 0 | 0 | 5 | 4 | 0 | 0 | 0 | 68 | 55 |
| New Jersey: | | | | | | | | | | | |
| Camden | 2 | | 0 | 2 | 4 | 12 | 0 | 2 | 0 | 1 | 36 |
| Newark | 1 | 4 | 2 | 3 | 7 | 10 | 0 | 7 | 1 | 25 | 100 |
| Trenton | 0 | | 1 | 0 | 5 | 8 | 0 | 0 | 0 | 0 | 40 |
| Pennsylvania: | | | | | | | | | | | |
| Philadelphia | 2 | 14 | 6 | 129 | 66 | 74 | 0 | 24 | 0 | 27 | 551 |
| Pittsburgh | 7 | 4 | 3 | 4 | 19 | 31 | 0 | 4 | 1 | 36 | 146 |
| Reading | 0 | | 0 | 1 | 5 | 1 | 0 | 0 | 0 | 15 | 33 |
| Ohio: | | | | | | | | | | | |
| Cincinnati | 14 | | 1 | 46 | 6 | 26 | 0 | 7 | 0 | 15 | 128 |
| Cleveland | 9 | 44 | 3 | 1 | 35 | 71 | 0 | 4 | 2 | 78 | 192 |
| Columbus | 5 | | 0 | 0 | 7 | 43 | 0 | 1 | 0 | 0 | 83 |
| Toledo | 2 | 1 | 1 | 34 | 6 | 38 | 0 | 7 | 0 | 9 | 80 |
| Indiana: | | | | | | | | | | | |
| Fort Wayne | 7 | | 0 | 0 | 2 | 8 | 0 | 1 | 0 | 0 | 23 |
| Indianapolis | 5 | | 0 | 1 | 8 | 7 | 0 | 0 | 0 | 18 | |
| South Bend | 0 | | 0 | 0 | 3 | 4 | 0 | 0 | 0 | 3 | 11 |
| Terre Haute | 3 | | 0 | 9 | 1 | 2 | 0 | 0 | 0 | 0 | 22 |
| Illinois: | | | | | | | | | | | |
| Chicago | 1 | 5 | 3 | 10 | 76 | 151 | 0 | 38 | 2 | 112 | 746 |
| Springfield | 2 | 2 | 0 | 0 | 2 | 6 | 0 | 0 | 0 | 1 | 20 |
| Michigan: | | | | | | | | | | | |
| Detroit | 9 | 4 | 5 | 20 | 32 | 62 | 0 | 8 | 0 | 82 | 269 |
| Flint | 0 | | 0 | 3 | 3 | 27 | 0 | 1 | 0 | 3 | 33 |
| Grand Rapids | 0 | | 1 | 0 | 1 | 4 | 0 | 0 | 0 | 0 | 25 |
| Wisconsin: | | | | | | | | | | | |
| Kenosha | 0 | | 0 | 0 | 2 | 17 | 0 | 0 | 0 | 4 | 15 |
| Madison | 0 | | | 1 | | 1 | 0 | | 0 | 33 | 6 |
| Milwaukee | 11 | | 0 | 5 | 8 | 20 | 1 | 8 | 0 | 62 | 86 |
| Racine | 0 | | 0 | 3 | 0 | 8 | 0 | 0 | 0 | 4 | 15 |
| Superior | 0 | | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 2 | 5 |
| Minnesota: | | | | | | | | | | | |
| Duluth | 0 | | 1 | 0 | 0 | 3 | 0 | 0 | 0 | 0 | 23 |
| Minneapolis | 3 | | 0 | 5 | 11 | 12 | 0 | 2 | 1 | 30 | 104 |
| St. Paul | 0 | 1 | 1 | 1 | 3 | 14 | 0 | 3 | 1 | 19 | 52 |
| Iowa: | | | | | | | | | | | |
| Des Moines | 2 | | | 0 | | 31 | 0 | | 0 | 0 | 21 |
| Sioux City | 2 | | | 0 | | 1 | 0 | | 0 | 3 | |
| Waterloo | 0 | | | 1 | | 0 | 0 | | 0 | 6 | |
| Missouri: | | | | | | | | | | | |
| Kansas City | 5 | | 1 | 2 | 7 | 30 | 0 | 5 | 0 | 6 | 139 |
| St. Joseph | 3 | | 0 | 0 | 3 | 2 | 0 | 0 | 0 | 0 | 18 |
| St. Louis | 22 | | | 59 | 12 | 23 | 0 | 8 | 4 | 33 | 185 |
| North Dakota: | | | | | | | | | | | |
| Fargo | 0 | | 0 | 8 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| Grand Forks | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 1 | 0 |
| South Dakota: | | | | | | | | | | | |
| Aberdeen | 0 | | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 |
| Sioux Falls | 0 | | 0 | 118 | 0 | 0 | 0 | 0 | 0 | 0 | 7 |
| Nebraska: | | | | | | | | | | | |
| Lincoln | 0 | | 0 | 1 | 0 | 4 | 1 | 0 | 0 | 1 | 0 |
| Omaha | 2 | | 0 | 8 | 2 | 14 | 1 | 0 | 0 | 8 | 52 |
| Kansas: | | | | | | | | | | | |
| Topeka | 0 | | 0 | 0 | 0 | 5 | 0 | 0 | 0 | 5 | 4 |
| Wichita | 0 | | 0 | 0 | 5 | 6 | 0 | 3 | 0 | 10 | 34 |
| Delaware: | | | | | | | | | | | |
| Wilmington | 0 | | 0 | 0 | 2 | 1 | 0 | 2 | 0 | 2 | 30 |
| Maryland: | | | | | | | | | | | |
| Baltimore | 6 | 16 | 3 | 2 | 25 | 22 | 0 | 7 | 1 | 53 | 227 |
| Cumberland | 1 | | 0 | 0 | 1 | 6 | 0 | 0 | 0 | 0 | 6 |
| Frederick | 0 | | 0 | 0 | 2 | 3 | 0 | 0 | 0 | 0 | 2 |
| District of Columbia: | | | | | | | | | | | |
| Washington | 9 | 2 | 0 | 31 | 13 | 17 | 0 | 13 | 2 | 20 | 167 |
| Virginia: | | | | | | | | | | | |
| Lynchburg | 5 | | 0 | 0 | 1 | 6 | 0 | 0 | 0 | 0 | 10 |
| Norfolk | 2 | | 0 | 1 | 0 | 9 | 0 | 0 | [1] 1 | 0 | 38 |
| Richmond | 7 | | 1 | 0 | 6 | 11 | 0 | 3 | 1 | 0 | 53 |
| Roanoke | 3 | | 0 | 0 | 1 | 6 | 0 | 0 | 0 | 0 | 11 |

[1] Nonresident.

*City reports for week ended Dec. 9, 1933*—Continued

| State and city | Diphtheria cases | Influenza | | Measles cases | Pneumonia deaths | Scarlet fever cases | Smallpox cases | Tuberculosis deaths | Typhoid fever cases | Whooping cough cases | Deaths all causes |
|---|---|---|---|---|---|---|---|---|---|---|---|
| | | Cases | Deaths | | | | | | | | |
| West Virginia: | | | | | | | | | | | |
| Charleston | 6 | | 0 | 0 | 3 | 2 | 0 | 1 | 0 | 0 | 27 |
| Huntington | 6 | | 0 | 0 | 0 | 14 | 0 | 0 | 0 | 0 | 0 |
| Wheeling | 0 | | 0 | 0 | 2 | 4 | 0 | 0 | 0 | 0 | 17 |
| North Carolina: | | | | | | | | | | | |
| Raleigh | 1 | | 0 | 0 | 1 | 2 | 0 | 0 | 0 | 1 | 23 |
| Wilmington | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 11 |
| Winston-Salem | 6 | | 0 | 153 | 4 | 8 | 0 | 2 | 0 | 0 | 20 |
| South Carolina: | | | | | | | | | | | |
| Charleston | 0 | 15 | 0 | 2 | 0 | 1 | 0 | 3 | 2 | 6 | 27 |
| Columbia | | | | | | | | | | | |
| Greenville | 1 | | 0 | 0 | 3 | 1 | 0 | 0 | 0 | 2 | 10 |
| Georgia: | | | | | | | | | | | |
| Atlanta | 12 | 24 | 1 | 3 | 10 | 6 | 0 | 3 | 0 | 3 | 79 |
| Brunswick | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 6 | 4 |
| Savannah | 0 | 27 | 0 | 0 | 1 | 2 | 0 | 3 | 3 | 0 | 29 |
| Florida: | | | | | | | | | | | |
| Miami | 1 | 1 | 1 | 0 | 0 | 1 | 0 | 2 | 0 | 0 | 30 |
| Tampa | 4 | 2 | 2 | 0 | 3 | 3 | 0 | 0 | 0 | 0 | 32 |
| Kentucky: | | | | | | | | | | | |
| Ashland | 2 | | | 0 | | 2 | 0 | | 0 | 0 | |
| Lexington | 2 | | 0 | 0 | 1 | 0 | 0 | 2 | 1 | 0 | 15 |
| Louisville | 14 | | 0 | 0 | 5 | 17 | 0 | 2 | 0 | 8 | 67 |
| Tennessee: | | | | | | | | | | | |
| Memphis | 6 | | 2 | 4 | 9 | 13 | 0 | 4 | 0 | 10 | 98 |
| Nashville | 1 | | 0 | 12 | 4 | 0 | 0 | 2 | 1 | 0 | |
| Alabama: | | | | | | | | | | | |
| Birmingham | 10 | 2 | 1 | 1 | 4 | 10 | 0 | 3 | 0 | 0 | 61 |
| Mobile | 3 | | 0 | 0 | 3 | 1 | 0 | 1 | 0 | 0 | 21 |
| Montgomery | 1 | 2 | | 0 | | 1 | 0 | | 0 | 0 | |
| Arkansas: | | | | | | | | | | | |
| Fort Smith | 2 | | | 0 | | 4 | 0 | | 0 | 2 | |
| Little Rock | 1 | | 0 | 3 | 6 | 6 | 0 | 1 | 0 | 0 | 7 |
| Louisiana: | | | | | | | | | | | |
| New Orleans | 10 | 11 | 5 | 0 | 20 | 11 | 0 | 11 | 4 | 0 | 160 |
| Shreveport | 1 | | 0 | 0 | 1 | 2 | 0 | 1 | 0 | 0 | 19 |
| Oklahoma: | | | | | | | | | | | |
| Tulsa | 1 | | | 7 | | 1 | 0 | | 0 | 0 | |
| Texas: | | | | | | | | | | | |
| Dallas | 21 | 3 | 3 | 0 | 14 | 5 | 1 | 3 | 2 | 4 | 72 |
| Fort Worth | 8 | | 1 | 0 | 9 | 15 | 0 | 0 | 0 | 0 | 39 |
| Galveston | 3 | | 0 | 0 | 3 | 3 | 0 | 1 | 1 | 0 | 20 |
| Houston | 29 | | 0 | 0 | 11 | 8 | 0 | 7 | 0 | 0 | 75 |
| San Antonio | 4 | | 1 | 0 | 6 | 6 | 0 | 1 | 1 | 0 | 57 |
| Montana: | | | | | | | | | | | |
| Billings | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 6 |
| Great Falls | 0 | | 0 | 0 | 0 | 1 | 0 | 0 | 1 | 2 | 10 |
| Helena | 0 | | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 5 |
| Missoula | 0 | | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 11 |
| Idaho: | | | | | | | | | | | |
| Boise | 0 | | 0 | 0 | 0 | 0 | 1 | 0 | 0 | 0 | 0 |
| Colorado: | | | | | | | | | | | |
| Denver | 3 | 37 | 1 | 3 | 11 | 9 | 0 | 4 | 0 | 37 | 92 |
| Pueblo | 0 | | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 10 | 7 |
| New Mexico: | | | | | | | | | | | |
| Albuquerque | 0 | | 0 | 0 | 1 | 2 | 0 | 2 | 0 | 1 | 10 |
| Utah: | | | | | | | | | | | |
| Salt Lake City | 0 | | 0 | 227 | 4 | 6 | 0 | 2 | 2 | 12 | 34 |
| Nevada: | | | | | | | | | | | |
| Reno | 0 | | 0 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 5 |
| Washington: | | | | | | | | | | | |
| Seattle | 0 | | | 0 | 8 | 5 | 0 | 5 | 0 | 54 | 80 |
| Spokane | 0 | 2 | 2 | 136 | 5 | 2 | 0 | 1 | 0 | 2 | 32 |
| Tacoma | 1 | | 0 | 0 | 2 | 2 | 0 | 1 | 0 | 7 | 25 |
| Oregon: | | | | | | | | | | | |
| Portland | 0 | | 0 | 2 | 6 | 20 | 0 | 3 | 0 | 2 | 78 |
| Salem | 0 | 2 | 0 | 1 | 0 | 0 | 0 | 0 | 0 | 6 | 0 |
| California: | | | | | | | | | | | |
| Los Angeles | 21 | 24 | 1 | 8 | 26 | 75 | 3 | 27 | 2 | 51 | 354 |
| Sacramento | 1 | | 0 | 16 | 6 | 1 | 0 | 2 | 0 | 1 | 26 |
| San Francisco | 2 | 4 | 2 | 0 | 10 | 9 | 0 | 9 | 0 | 9 | 195 |

*City reports for week ended Dec. 9, 1933*—Continued

| State and city | Meningococcus meningitis | | Poliomyelitis cases | State and city | Meningococcus meningitis | | Poliomyelitis cases |
|---|---|---|---|---|---|---|---|
| | Cases | Deaths | | | Cases | Deaths | |
| Massachusetts: | | | | Kansas: | | | |
| Fall River | 0 | 0 | 1 | Topeka | 1 | 0 | 0 |
| New York: | | | | West Virginia: | | | |
| New York | 7 | 4 | 2 | Wheeling | 1 | 0 | 0 |
| Pennsylvania: | | | | North Carolina: | | | |
| Philadelphia | 0 | 1 | 0 | Wilmington | 0 | 1 | 0 |
| Ohio: | | | | Georgia: | | | |
| Cleveland | 1 | 1 | 2 | Atlanta | 1 | 1 | 0 |
| Toledo | 1 | 0 | 0 | Washington: | | | |
| Indiana: | | | | Seattle | 0 | 0 | 2 |
| Indianapolis | 2 | 0 | 0 | California: | | | |
| Illinois: | | | | Los Angeles | 1 | 0 | 0 |
| Chicago | 1 | 1 | 0 | San Francisco | 0 | 0 | 1 |
| Iowa: | | | | | | | |
| Des Moines | 0 | 0 | 1 | | | | |

*Lethargic encephalitis.*—Cases: New York, 1; Chicago, 1; St. Louis, 5; Topeka, 1; Richmond, Va., 1; Birmingham, 1; Dallas, Tex., 1; Portland, Oreg., 1.

*Typhus fever.*—Cases: New York, 1; Charleston, S.C., 1; Atlanta, 3; Savannah, 3; Dallas, 1; Fort Worth, Tex., 1.

*Pellagra.*—Cases: Baltimore, 1; Washington, 1.

# FOREIGN AND INSULAR

## CANADA

*Provinces—Communicable diseases—2 weeks ended December 2, 1933.*—During the 2 weeks ended December 2, 1933, cases of certain communicable diseases were reported by the Department of Pensions and National Health of Canada, as follows:

| Disease | Prince Edward Island | Nova Scotia | New Brunswick | Quebec | Ontario | Manitoba | Saskatchewan | Alberta [1] | British Columbia | Total |
|---|---|---|---|---|---|---|---|---|---|---|
| Cerebrospinal meningitis | | | | | 1 | 1 | 1 | 1 | 2 | 6 |
| Chicken pox | | 6 | 3 | 423 | 550 | 166 | 128 | 26 | 152 | 1,454 |
| Diphtheria | | 3 | 1 | 56 | 39 | 33 | 1 | 1 | | 134 |
| Dysentery | | | | 1 | 1 | | 1 | | 1 | 4 |
| Erysipelas | | | | 9 | 7 | 2 | 1 | | 1 | 20 |
| Influenza | | 15 | | 13 | 11 | | | | 39 | 78 |
| Lethargic encephalitis | | | | | | | 1 | | | 1 |
| Measles | | | | 112 | 22 | | 34 | 1 | 7 | 176 |
| Mumps | | | | | 130 | 7 | 1 | 1 | 78 | 217 |
| Paratyphoid fever | 1 | | | | | | | | | 1 |
| Pneumonia | | 1 | | | 36 | | 5 | | 11 | 53 |
| Poliomyelitis | | | | 1 | 1 | | | | | 2 |
| Scarlet fever | | 23 | 12 | 163 | 231 | 80 | 12 | 1 | 130 | 652 |
| Smallpox | | | | 1 | | | | | | 1 |
| Trachoma | | | | 1 | 1 | 2 | | | | 4 |
| Tuberculosis | 5 | 2 | 10 | 129 | 77 | 44 | 6 | | 47 | 320 |
| Typhoid fever | | | 4 | 53 | 21 | | 1 | | 2 | 81 |
| Whooping cough | | 11 | 3 | 259 | 120 | 154 | 41 | 5 | 23 | 616 |

[1] No report was received from Alberta for the week ended Dec. 2, 1933.

*Ontario Province—Communicable diseases—Four weeks ended November 25, 1933.*—The Department of Health of the Province of Ontario, Canada, reports certain communicable diseases for the 4 weeks ended November 25, 1933, as follows:

| Disease | Cases | Deaths | Disease | Cases | Deaths |
|---|---|---|---|---|---|
| Cerebrospinal meningitis | 3 | 1 | Pneumonia | | 138 |
| Chicken pox | 908 | | Poliomyelitis | 3 | |
| Diphtheria | 56 | 3 | Puerperal septicemia | | 1 |
| Dysentery | 1 | | Scarlet fever | 449 | |
| Erysipelas | 13 | 1 | Septic sore throat | 5 | |
| German measles | 14 | | Syphilis | 81 | 1 |
| Gonorrhea | 116 | | Tetanus | 1 | |
| Influenza | 15 | 2 | Tuberculosis | 164 | 26 |
| Lethargic encephalitis | 1 | | Tularaemia | 1 | |
| Measles | 77 | | Typhoid fever | 45 | 1 |
| Mumps | 175 | | Undulant fever | 6 | |
| Paratyphoid fever | 7 | | Whooping cough | 285 | 8 |

## JAMAICA

*Communicable diseases—Four weeks ended December 2, 1933.*—During the 4 weeks ended December 2, 1933, cases of certain communicable diseases were reported in Kingston, Jamaica, and in the island outside of Kingston, as follows:

| Disease | Kingston | Other localities | Disease | Kingston | Other localities |
|---|---|---|---|---|---|
| Chicken pox | 2 | 9 | Lethargic encephalitis | | 2 |
| Diphtheria | | 2 | Puerperal fever | | 2 |
| Dysentery | 9 | 15 | Tuberculosis | 38 | 77 |
| Erysipelas | | 3 | Typhoid fever | 16 | 74 |
| Leprosy | | 1 | | | |

## CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER

From medical officers of the Public Health Service, American consuls, International Office of Public Hygiene, Pan American Sanitary Bureau, health section of the League of Nations, and other sources. The reports contained in the following tables must not be considered as complete or final as regards either the list of countries included or the figures for the particular countries for which reports are given.

### CHOLERA

[C indicates cases; D, deaths; P, present]

| Place | | Apr. 30–May 27, 1933 | May 28–June 24, 1933 | June 25–July 29, 1933 | July 30–Aug. 26, 1933 | Week ended— September 1933 2 | 9 | 16 | 23 | 30 | October 1933 7 | 14 | 21 | 28 | November 1933 4 | 11 | 18 | 25 | Dec. 2, 1933 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| China: | | | | | | | | | | | | | | | | | | | |
| Canton | C | | | 1 | | | | | | | | | | | | | | | |
| Hankow | C | 2 | 1 | 2 | | | 1 | | | | | | | | | | | | |
| | D | 1 | | 1 | 1 | | | | | | | | | | | | | | |
| Swatow | C | 2 | | | | | | | | | | | | | | | | | |
| Tientsin | C | | | 4 | | | | | | | | | | | | | | | |
| India | C | 8,201 | 3,517 | 6,881 | 7,695 | 2,901 | 2,576 | 2,534 | 3,427 | 2,984 | 2,211 | 2,091 | | | | | | | |
| | D | 4,091 | 1,555 | 3,047 | 3,237 | 1,315 | 1,166 | 1,233 | 1,502 | 1,438 | 1,097 | 943 | | | | | | | |
| Bombay Presidency | C | | | 3,869 | 3,604 | 651 | 610 | 532 | 642 | 646 | 407 | | 421 | 548 | 352 | | | | |
| | D | | | 1,609 | 1,436 | 305 | 282 | 259 | 301 | 265 | 199 | | 169 | 266 | 175 | | | | |
| Bombay | C | 2 | 1 | 6 | 1 | | | | 1 | 4 | 1 | | | 3 | 1 | | | | 4 |
| Calcutta | C | 690 | 196 | 77 | 59 | 21 | 28 | 23 | 12 | 9 | 26 | 15 | 11 | 16 | 13 | 11 | | 18 | 10 |
| Chittagong | C | 21 | 7 | | 1 | | | | | | | 1 | | | | | | | 1 |
| Madras Presidency | C | | | 276 | 923 | 274 | 213 | 182 | 247 | 142 | 142 | 90 | 155 | | | | | | |
| | D | | | 74 | 338 | 126 | 94 | 79 | 91 | 54 | 46 | 51 | 68 | | | | | | |
| Madras | C | | | | | | | | | | | | | 4 | 1 | | 2 | 3 | 61 |
| | D | | | | | | | | | | | | | 1 | 2 | | | 2 | 26 |
| Moulmein | C | | 3 | | | | | | | | | | | | | | | | |
| Rangoon | C | 1 | 1 | 2 | | | | | | | | | | | | | | | |
| Vizagapatam | C | 1 | | 4 | | | | | | | | | 1 | | | | | | |
| India (French): | | | | | | | | | | | | | | | | | | | |
| Chandernagor | D | | | | | | | | | 2 | | | | | | | | | |
| Karikal | C | | | | | | | | | 1 | | | | | | | | | |
| | D | | | | | | | | | 1 | | | | | 4 | | | | |
| Pondichery | C | | | | | 1 | | | | 1 | | | | | | | | | |
| | D | | | | | 1 | | | | 1 | | | | | | | | | |
| Indo-China (see also table below): | | | | | | | | | | | | | | | | | | | |
| Pnom-Penh | C | 2 | 1 | | | | | | | | | | | | | | | | |
| Saigon and Cholon | C | | | 8 | 2 | | | | | | | | | | | | | | |

## CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

### CHOLERA—Continued

[C indicates cases; D, deaths; P, present]

| Place | | Apr. 30–May 28, 1933 | May 28–June 24, 1933 | June 25–July 29, 1933 | July 30–Aug. 26, 1933 | Week ended— | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | | September 1933 | | | | | October 1933 | | | | November 1933 | | | | Dec. 2, 1933 |
| | | | | | | 2 | 9 | 16 | 23 | 30 | 7 | 14 | 21 | 28 | 4 | 11 | 18 | 25 | |
| Philippine Islands: [1] | | | | | | | | | | | | | | | | | | | |
| Antique Province | C | | | | | | | | | | | | 9 | 8 | 2 | | | 6 | 8 |
| | D | | | | | | | | | | | | 6 | 10 | 1 | | | 5 | 6 |
| Bohol Province | C | 15 | 80 | | 1 | | | | | 6 | 1 | 28 | 6 | 52 | 35 | 31 | 39 | 68 | 51 |
| | D | 12 | 38 | | | | | | | 3 | 1 | 12 | 3 | 31 | 21 | 14 | 25 | 41 | 25 |
| Cebu Province | C | | 2 | 90 | 26 | 1 | 9 | 6 | 11 | 12 | 22 | 16 | 15 | 10 | 23 | 18 | 7 | 23 | 15 |
| | D | | 2 | 39 | 20 | 1 | 7 | 4 | 9 | 7 | 14 | 9 | 9 | 4 | 16 | 12 | 4 | 13 | 10 |
| Cebu | C | | 1 | | 1 | 1 | | 1 | 2 | 5 | 6 | 6 | 3 | 6 | 1 | 11 | 4 | 1 | |
| | D | | 1 | | 1 | 1 | | 1 | 2 | 3 | 2 | 3 | 1 | 2 | 1 | 6 | 1 | | |
| Naga | C | | | | | | | | | | | | | | 9 | 1 | 1 | 7 | 4 |
| | D | | | | | | | | | | | | | | 6 | 1 | 1 | 1 | 3 |
| Iloilo Province | C | 1 | | 1 | | | | | | | | | | 12 | 1 | | | 1 | |
| | D | 1 | | 1 | | | | | | | | | | 11 | 2 | | | 1 | |
| Iloilo | C | 1 | | 1 | | | | | | | | | | | 1 | | | 1 | |
| | D | 1 | | 1 | | | | | | | | | | | 2 | | | 1 | |
| Leyte Province | C | 22 | 2 | | 2 | | 1 | | | | | | | | | | | | |
| | D | 20 | 2 | | 2 | | 1 | | | | | | | | | | | | |
| Occidental Negros Province | C | | 19 | 6 | 1 | | | | | | | | | | | | | | 2 |
| | D | | 9 | 7 | 1 | | | | | | | | | | | | | | 1 |
| Pampanga Province | C | 1 | | | | | | | | | | | | | | | | | |
| | D | 1 | | | | | | | | | | | | | | | | | |
| Rizal Province | C | 1 | | | | | | | | | | | | | | | | | |
| Samar Province | C | 29 | 7 | | 31 | | | | | [2] 31 | 11 | | | | | | | | |
| | D | 23 | 5 | | 20 | | | | | [2] 20 | 8 | | | | | | | | |
| Siam: | | | | | | | | | | | | | | | | | | | |
| Bangkok | C | 1 | 1 | | | | | | | | | | | | | | | | |
| Provinces | C | 13 | 28 | 1 | | | | | | | | | | | | | | | |
| | D | 6 | 14 | 1 | | | | | | | | | | | | | | | |
| On vessels: | | | | | | | | | | | | | | | | | | | |
| S.S. Dunana at Madras | D | 1 | | | | | | | | | | | | | | | | | |
| S.S. Lauterfels at Calcutta | C | 1 | | | | | | | | | | | | | | | | | |
| S.S. Mathura at Calcutta | C | 1 | | | | | | | | | | | | | | | | | |

| Place | | May 1933 | | | June 1933 | | | July 1933 | | | August 1933 | | | September 1933 | | | October 1933 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | 1-10 | 11-20 | 21-31 | 1-10 | 11-20 | 21-30 | 1-10 | 11-20 | 21-31 | 1-10 | 11-20 | 21-31 | 1-10 | 11-20 | 21-30 | 1-10 | 11-20 |
| Indo-China (French) (see also table above): | | | | | | | | | | | | | | | | | | |
| Cambodia [3] | C | 11 | 14 | 17 | 23 | 31 | 14 | 3 | 1 | ---- | ---- | 1 | ---- | ---- | ---- | ---- | ---- | ---- |
| | D | 8 | 10 | 9 | 12 | 17 | 6 | 3 | 1 | ---- | ---- | 1 | ---- | ---- | ---- | ---- | ---- | ---- |
| Cochin-China [3] | C | 5 | 9 | 6 | 4 | 8 | 5 | 6 | 2 | 12 | 3 | 2 | 1 | 1 | 2 | 2 | 5 | 5 |
| | D | 4 | 8 | 6 | 4 | 8 | 4 | 5 | 2 | 3 | 3 | 2 | 1 | 1 | 2 | 2 | 5 | 3 |

[1] During the week ended Dec. 16, 1933, cholera was reported in the Philippine Islands as follows: Bohol Province—Calape, 3 cases, 3 deaths; Loon, 8 cases, 5 deaths; Tubigon, 11 cases, 8 deaths. Cebu Province—Liloan, 1 case, 1 death. Oriental Negros Province—Tanjay, 4 cases, 4 deaths.
[2] For 2 weeks.
[3] Reports incomplete.

## PLAGUE [1]

[C indicates cases; D, deaths; P, present]

| Place | | Apr. 30-May 27, 1933 | May 28-June 24, 1933 | June 25-July 29, 1933 | July 30-Aug. 26, 1933 | Week ended— | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | | September 1933 | | | | | October 1933 | | | | November 1933 | | | |
| | | | | | | 2 | 9 | 16 | 23 | 30 | 7 | 14 | 21 | 28 | 4 | 11 | 18 | 25 |
| Argentina (see also table below): Cordoba Province | C | 8 | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- |
| Azores: | | | | | | | | | | | | | | | | | | |
| Fayal | C | ---- | ---- | ---- | ---- | ---- | ---- | ---- | 1 | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- |
| St. Michaels | C | ---- | ---- | ---- | ---- | ---- | ---- | ---- | 7 | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- |
| | D | ---- | ---- | ---- | ---- | ---- | ---- | ---- | 1 | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- |
| Bolivia. (See table below.) | | | | | | | | | | | | | | | | | | |
| British East Africa (see also table below): | | | | | | | | | | | | | | | | | | |
| Kenya | C | ---- | 3 | 7 | 13 | 2 | 6 | 3 | 10 | 9 | 16 | 2 | ---- | ---- | 8 | ---- | ---- | ---- |
| Tanganyika | D | 1 | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | 1 | ---- | ---- | ---- |
| Uganda | C | 85 | 53 | 58 | 77 | 18 | 34 | 22 | 21 | 18 | 8 | 22 | 15 | ---- | ---- | ---- | ---- | ---- |
| | D | 83 | 53 | 55 | 77 | 18 | 28 | 23 | 23 | 17 | 7 | 22 | 14 | ---- | ---- | ---- | ---- | ---- |
| Ceylon: Colombo | C | ---- | 1 | 2 | ---- | ---- | 1 | ---- | ---- | ---- | ---- | ---- | 1 | ---- | ---- | ---- | ---- | ---- |
| | D | ---- | 1 | 2 | ---- | ---- | 1 | ---- | ---- | ---- | ---- | ---- | 1 | ---- | ---- | ---- | ---- | ---- |
| Plague-infected rats | | ---- | ---- | 4 | 1 | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- | ---- |
| China: Manchuria.[2] | | | | | | | | | | | | | | | | | | |

See footnotes at end of table.

## CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

### PLAGUE—Continued

[C indicates cases; D, deaths; P, present]

| Place | | Apr. 30–May 27, 1933 | May 28–June 24, 1933 | June 25–July 29, 1933 | July 30–Aug. 26, 1933 | Week ended— September 1933 2 | 9 | 16 | 23 | 30 | October 1933 7 | 14 | 21 | 28 | November 1933 4 | 11 | 18 | 25 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Dutch East Indies: West Java | C | 878 | 793 | 1,434 | 899 | 248 | 299 | 289 | 294 | 335 | | | | | | | | |
| | D | 872 | 789 | 1,428 | 898 | 248 | 299 | 288 | 294 | 334 | | | | | | | | |
| Ecuador. (See table below.) | | | | | | | | | | | | | | | | | | |
| Egypt: | | | | | | | | | | | | | | | | | | |
| Alexandria | C | | | 4 | 1 | | | | | | | | | | | | | |
| Asyut | C | 10 | 2 | | | | | | | | | | | | | | | |
| Aswan | C | 1 | | | | | | | | | | | | | | | | |
| Faiyum | C | | 2 | 4 | | | 1 | 1 | | | 1 | | | | | | | |
| | D | | | 1 | | | | | | | | | | | | | | |
| Gharbiya | C | | | | | | | | | 3 | 1 | | | 1 | | | | |
| | D | | | | | | | | | 2 | 1 | 1 | | 1 | | | | |
| Girga | C | | | 2 | | | | | | | | | | | | | | |
| Minufiya | C | 1 | | | | | | | | | | | | | | | | |
| Minya | C | | 4 | 1 | | | | | | 1 | | | | | | | | |
| Qena | C | | | | 1 | | | | | | | | | | | | | |
| | D | | | | 1 | | | | | | | | | | | | | |
| Luxor | C | | | | 2 | | | | | | | | | | | | | |
| France: Marseille | C | | | | 8 | | | | | | | | | | | | | |
| | D | | | | 3 | | | | | | | | | | | | | |
| Plague-infected rats | | | | | 2 | | | | | | | | | | | 1 | | |
| Hawaii Territory: Hawaii Island—Hamakua—Plague-infected rats | | | | | | | 1 | | | | | | | | | 2 | | |
| India | C | 2,319 | 1,411 | 3,869 | 6,209 | 2,151 | 2,350 | 2,610 | 3,133 | 3,398 | 3,074 | 2,941 | | | | | | |
| | D | 1,407 | 1,231 | 2,616 | 3,560 | 1,331 | 1,369 | 1,573 | 1,747 | 1,951 | 1,603 | 1,546 | | | | | | |
| Bassein | C | 3 | 1 | 11 | 7 | | | 2 | | 1 | | | | | | | | |
| Plague-infected rats | | 3 | 2 | 2 | 1 | | | | | | | | | | | | | |
| Bombay Presidency | C | | | 2,448 | 3,971 | 1,263 | 1,424 | 1,465 | 1,872 | 2,045 | 2,231 | | 1,382 | 1,309 | 1,598 | | | |
| | D | | | 1,493 | 2,313 | 852 | 898 | 967 | 1,173 | 1,227 | 1,235 | | 895 | 708 | 992 | | | |
| Bombay | C | 6 | 2 | 1 | 1 | 2 | 1 | 1 | 2 | 1 | 1 | 1 | | 1 | | 1 | | |
| Plague-infected rats | | 34 | 16 | 5 | 5 | | | 1 | | 2 | 1 | | | | 3 | 3 | 1 | 1 |
| Poona | C | | | | | | | | | | 176 | | 150 | 149 | 61 | | | |
| | D | | | | | | | | | | 118 | 199 | 109 | 101 | 53 | | | |
| Madras Presidency | C | 23 | 8 | 372 | 867 | 253 | 190 | 183 | 273 | 282 | 231 | 100 | 95 | | | | | |
| | D | 8 | 6 | 148 | 395 | 136 | 77 | 83 | 111 | 140 | 139 | 56 | 46 | | | | | |
| Rangoon | C | 1 | 1 | | | | | 2 | | | 1 | | | | | | | 1 |
| Plague-infected rats | | | | 3 | 1 | | | | | 2 | | | | | | | 1 | 1 |

| Place | | May 1933 | June 1933 | July 1933 | August 1933 | September 1933 | October 1933 |
|---|---|---|---|---|---|---|---|
| Indo-China (see also table below): | | | | | | | |
| Pnom-Penh | D | 6 | 3 | 2 | 3 | 2 | |
| Saigon and Cholon | C | 2 | | 2 | 3 | | |
| Plague-infected rats | | 3 | | | | | |
| Iraq: | | | | | | | |
| Baghdad | C | 3 | 2 | 7 | | 1 | 1 |
| Basra | C | | | 3 | | | |
| Libya: Gheran | C | | | | | | |
| | D | | | | | | |
| Madagascar (see also table below): Tamatave | C | | | | 1 | | |
| Morocco | C | 2 | | 8 | | | |
| Peru. (See table below.) | | | | | | | |
| Senegal. (See table below.) | | | | | | | |
| Siam | C | | 2 | | | | |
| South-West Africa.[4] | | | | | | | |
| Syria: Beirut | C | | 1 | 2 | | | |
| Union of South Africa: Orange Free State | C | 7 | | | | | |
| United States: California: | | | | | | | |
| San Benito County—Plague-infected ground squirrels | | | | | 8 | | |
| Whittier | C | | | | 1 | | |
| | D | | | | 1 | | |
| On vessel: S.S. Angkor at Beirut from Marseille | C | | | | | | |

| Place | | May 1933 | June 1933 | July 1933 | August 1933 | September 1933 | October 1933 |
|---|---|---|---|---|---|---|---|
| Argentina (see also table above) | C | | | 7 | | | 6 |
| | D | | | 3 | | | 2 |
| Bolivia | C | | | 25 | 1 | | |
| British East Africa (see also table above): | | | | | | | |
| Kenya | C | 4 | 6 | 3 | 13 | 36 | 20 |
| Uganda | C | | | 47 | 91 | 97 | 71 |
| Ecuador | C | 35 | | | | 3 | |
| Indo-China (see also table above): | | | | | | | |
| Cambodia | D | 3 | 5 | 3 | 6 | 16 | 8 |
| Cochin-China | C | 2 | | 2 | 5 | 1 | |
| Madagascar | C | | | 133 | 100 | | |
| | D | | | 132 | 92 | | |
| Peru | C | 3 | 5 | 6 | 7 | 8 | 18 |
| Callao | C | | | | | 1 | 7 |
| Senegal: | | | | | | | |
| Dakar [5] | C | 2 | | 57 | | 5 | 4 |
| | D | 1 | | 39 | | 3 | 3 |
| Tivaouane [5] | C | | 1 | | 5 | 2 | |

[1] Including plague in the United States and its possessions.
[2] In September and October 1933 plague was reported in parts of Manchuria, China, especially between the Ssupingkai-Taonan Railway and the southern line of the Chinese Eastern Railway, also adjacent to the lines of the Ssupingkai-Taonan, Ssupingkai-Tungliao, and Tahushan-Tungliao Railways.
[3] Imported.
[4] 103 cases of plague with 5 deaths were reported in Ovamboland, South-West Africa from Jan. 1 to Oct. 14, 1933. Antiplague measures have been taken.
[5] Incomplete reports.

20288°—34——3

## CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

### SMALLPOX

[C indicates cases; D, deaths; P, present]

| Place | | Apr. 30–May 27, 1933 | May 28–June 24, 1933 | June 25–July 29, 1933 | July 30–Aug. 26, 1933 | Week ended— | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | | September 1933 | | | | | October 1933 | | | | November 1933 | | | | Dec. 2, 1933 |
| | | | | | | 2 | 9 | 16 | 23 | 30 | 7 | 14 | 21 | 28 | 4 | 11 | 18 | 25 | |
| Algeria: | | | | | | | | | | | | | | | | | | | |
| Algiers Department | C | | | 2 | | | | | | | | | | | | 1 | | | |
| Constantine Department | C | | | 1 | | | | | | | | | | | | | | | |
| Arabia: Muscat—Oman Sultanate | C | | 73 | | | | | | | | | | | | | | | | |
| | D | | 20 | | | | | | | | | | | | | | | | |
| Belgian Congo | C | | | 5 | | 2 | 2 | | | | | | | | | | | | |
| | D | | | 1 | | | | | | | | | | | | | | | |
| Bolivia. (See table below.) | | | | | | | | | | | | | | | | | | | |
| Brazil: Porto Alegre (alastrim) | C | | 6 | 1 | 1 | | 1 | | 1 | 10 | | | 1 | | | | | | |
| British East Africa: | | | | | | | | | | | | | | | | | | | |
| Kenya | C | 4 | | | | | | | | | | | | | | | | | |
| Tanganyika | C | 13 | 10 | 6 | 21 | 9 | | 16 | 2 | 3 | 12 | 12 | 24 | 4 | | | | | |
| British South Africa: | | | | | | | | | | | | | | | | | | | |
| Northern Rhodesia | C | 2 | | 1 | 1 | | | | | | | | | | | | | | |
| Southern Rhodesia | C | 16 | | 1 | | | | 2 | | | | | | | | | | | |
| Canada: | | | | | | | | | | | | | | | | | | | |
| British Columbia | C | [1] 1 | [1] 1 | | | | | | | | | | | | | | | | |
| Manitoba | C | | | | 1 | | | | | | | | | | | | | | |
| Ontario | C | [1] 1 | [1] 1 | | | | | | | | | | | | | | | | |
| Toronto | C | 1 | | | | | | | | | | | | | | | | | |
| Saskatchewan | C | [1] 5 | 7 | | | | [1] 1 | | | | | | | | | | | | |
| Ceylon: | | | | | | | | | | | | | | | | | | | |
| Colombo | C | 4 | 9 | 6 | 9 | | | | | | | | | | | 1 | | | |
| Galle | C | 2 | | 1 | | | | | | | | | | | | | | | |
| China: | | | | | | | | | | | | | | | | | | | |
| Amoy | C | 3 | 2 | | | | | | | | | | | | | | | | |
| Canton | C | 7 | | | | | | | | | | | | | 1 | | 1 | | 1 |
| Dairen | C | 3 | 2 | 3 | | | 1 | | | | | 1 | 1 | 1 | 2 | | | | |
| Foochow | C | P | P | P | | | | | | | | | | | | | | | |
| Hangchow | C | | | 3 | 1 | | | | | | | | | | | | | | |
| | D | | | 2 | 1 | | | | | | | | | | | | | | |
| Hong Kong | C | 19 | 8 | 3 | 1 | | | | | | | | | | | | | | |
| Macao | C | 4 | | | | | | | | | | | | | | | | | |
| Nanking | C | 2 | 2 | 2 | | | | | | | | | | | | | | | |
| Shanghai | C | 33 | 27 | 22 | 10 | 2 | | 1 | 1 | 1 | 2 | | | 1 | 4 | 1 | | 6 | |
| Swatow | C | 10 | 4 | | | | | | | | | 1 | 1 | | 1 | | | | |
| Tientsin | C | | 16 | 2 | | | | | | | | | | | | | | | |

| | | | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Chosen. (See table below.) | | | | | | | | | | | | | | | | | | | |
| Colombia: Buenaventura | D | | | 1 | | | | | | | | | | | | | | | |
| Costa Rica. (See table below.) | | | | | | | | | | | | | | | | | | | |
| Dahomey. (See table below.) | | | | | | | | | | | | | | | | | | | |
| Ecuador. (See table below.) | | | | | | | | | | | | | | | | | | | |
| Egypt: | | | | | | | | | | | | | | | | | | | |
| Alexandria | C | 11 | | | | | | | | | | | | | 1 | 2 | | | |
| Asyut | C | | | 112 | 109 | | | 17 | | 26 | 6 | 8 | 21 | 5 | | 1 | | | |
| | D | | | 23 | 11 | | | 3 | | 3 | 1 | 1 | 1 | 1 | | | | | |
| Beheira | C | 62 | 6 | | | | | | | | | | | | | | | | |
| Beni-Suef | C | | | | | | | 10 | 17 | 4 | 10 | 2 | 1 | | | | | | |
| | D | | | | | | | 4 | 1 | 2 | 1 | 1 | | | | | | | |
| Cairo | C | 10 | 4 | 1 | | | | | | 1 | | | | | | | | | |
| Dakahliya | C | | | 4 | 1 | | | | | | | | | | | | | | |
| Faiyum | C | | | 44 | 18 | | | 2 | | 1 | 1 | | | | | | | | |
| Gharbiya | C | 93 | 142 | 83 | 39 | 4 | | 1 | | | | 1 | | 1 | | 4 | | | |
| Girga | C | | | 2 | 2 | | | | | | | 1 | 5 | 3 | | | | | |
| Minufiya | C | 128 | 26 | | | | | | | | | | | | | | | | |
| Qena | C | | | 9 | 25 | | | 3 | 1 | | | | 13 | 12 | | 6 | | | |
| Provinces | C | 347 | 287 | 499 | 297 | 50 | 32 | 33 | | 35 | 18 | 12 | 40 | 25 | 25 | | | | |
| France. (See table below.) | | | | | | | | | | | | | | | | | | | |
| Great Britain: | | | | | | | | | | | | | | | | | | | |
| England and Wales | C | 91 | 71 | 55 | 32 | 4 | 2 | 6 | 4 | 4 | 3 | 3 | | 4 | 1 | 1 | 1 | 6 | |
| London | C | 79 | 51 | 40 | 28 | 3 | | 5 | 3 | 3 | 3 | 3 | | 4 | 1 | 1 | 1 | 6 | |
| London and Great Towns | C | 82 | 56 | 49 | 31 | 3 | 1 | 5 | 3 | 3 | 3 | 3 | | 4 | 1 | 1 | 1 | 6 | |
| Greece (see also table below): Salonika | C | 18 | | 8 | | | | | 9 | | | | | | | | | | |
| Guatemala. (See table below.) | | | | | | | | | | | | | | | | | | | |
| Honduras: Tegucigalpa | C | 7 | 3 | | | | | | | | | | | | | | | | |
| India | C | 33,876 | 23,631 | 14,599 | 7,069 | 1,765 | 1,509 | 1,156 | 1,377 | 1,245 | 1,136 | 1,107 | | | | | | | |
| | D | 9,706 | 7,008 | 4,256 | 1,768 | 430 | 345 | 268 | 321 | 261 | 310 | 215 | | | | | | | |
| Bassein | C | 32 | 16 | 9 | 10 | 4 | 1 | 4 | 7 | 3 | 3 | 3 | 2 | 4 | 5 | 3 | 2 | 2 | 4 |
| | D | 11 | 11 | 8 | 6 | 3 | 1 | | 1 | 1 | 3 | | 2 | 2 | 1 | 1 | 3 | 1 | 3 |
| Bombay Presidency | C | | | 1,942 | 1,029 | 198 | 170 | 174 | 202 | 129 | 130 | 191 | 127 | 151 | 121 | | | | |
| | D | | | 376 | 186 | 40 | 38 | 20 | 46 | 16 | 23 | 14 | 29 | 25 | 22 | | | | |
| Bombay | C | 212 | 88 | 34 | 14 | 1 | | | | | 2 | | 1 | 1 | | 2 | | 2 | 3 |
| | D | 148 | 58 | 25 | 8 | 1 | | | | | 2 | | 1 | 1 | | 2 | | 1 | 3 |
| Calcutta | C | 342 | 134 | 59 | 18 | 4 | 2 | 2 | 2 | | 1 | 3 | | 4 | 1 | 8 | | 4 | 9 |
| | D | 289 | 107 | 49 | 8 | 2 | | | 1 | | 1 | 1 | | 1 | 1 | 1 | | 3 | 1 |
| Chittagong | C | 1 | | | | | | | | | | | | | | | | | |
| | D | 1 | | | | | | | | | | | | | | | | | |
| Cochin | C | | 2 | 4 | | 2 | | | | | | | | 1 | | | | | |
| Karachi | C | 52 | 19 | 16 | 5 | | | | | | | | 2 | | | | | | |
| Madras Presidency | C | | | 3,571 | 2,851 | 621 | 592 | 539 | 548 | 644 | 602 | 540 | 646 | | | | | | |
| | D | | | 726 | 587 | 114 | 93 | 102 | 100 | 104 | 132 | 106 | 142 | | | | | | |
| Madras | C | 231 | 144 | 234 | 223 | 63 | 33 | 32 | 36 | 45 | 30 | 22 | 13 | 16 | 13 | 7 | 7 | 14 | 14 |
| Moulmein | C | | 12 | | | | | | | | | | | | | | | | |
| Negapatam | C | 7 | 3 | 12 | 21 | 2 | 2 | 3 | | 1 | | 1 | 4 | 4 | 7 | | 3 | 7 | 8 |
| Rangoon | C | 17 | 6 | 4 | 1 | 1 | | 3 | | | 1 | | | | | 1 | | 1 | 2 |
| Tuticorin | C | | | | 2 | | | | | | | | | 1 | 5 | | | 1 | |
| Vizagapatam | C | 2 | 5 | 14 | 11 | | | | 2 | | | 1 | | | 1 | | | 1 | |

[1] For 2 weeks.

## CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

### SMALLPOX—Continued

[C indicates cases; D, deaths; P, present]

| Place | | Apr. 30–May 27, 1933 | May 28–June 24, 1933 | June 25–July 29, 1933 | July 30–Aug. 26, 1933 | Week ended— | | | | | | | | | | | | Dec. 2, 1933 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | | September 1933 | | | | | October 1933 | | | | November 1933 | | | |
| | | | | | | 2 | 9 | 16 | 23 | 30 | 7 | 14 | 21 | 28 | 4 | 11 | 18 | 25 |
| India (French): | | | | | | | | | | | | | | | | | | |
| Chandernagor | D | | | 1 | | | | | | | | | | | | | | |
| Karikal | C | | 5 | | | | | | | 1 | 1 | | | | | | | |
| | D | | 5 | | | | | | | 1 | 1 | | | | | | | |
| Pondichery | C | | 129 | 139 | 181 | 57 | 50 | 40 | 24 | 39 | 59 | 25 | | 17 | 15 | | | |
| | D | | 82 | 73 | 70 | 22 | 25 | 17 | 22 | 18 | 32 | 19 | | 12 | 7 | | | |
| Indo-China (see also table below): Saigon and Cholon | C | | | 1 | | | 1 | | | | | | | | | | | |
| Iraq: | | | | | | | | | | | | | | | | | | |
| Baghdad | C | 5 | | | | | | | | | 1 | | 1 | | | | 1 | 1 |
| Basra | C | 6 | 2 | | | 4 | | | | | | | | | 3 | | | |
| Japan: | | | | | | | | | | | | | | | | | | |
| Kobe | C | | | 1 | | | | | | | | | | | | | | |
| Osaka | C | 1 | | | | | | | | | | | | | | | | |
| Tokyo | C | 1 | 2 | 45 | | | | | | | | | | | | | | |
| Mexico (see also table below): | | | | | | | | | | | | | | | | | | |
| Coahuila | C | 1 | 2 | | | | | | | | | | | | | | | |
| Juarez [2] | C | 2 | | | | | | | | | | | | | | | | |
| Mexico, D.F. | C | 5 | 5 | 1 | 12 | | | | | | | | | | | | | |
| Monterrey | D | 2 | | | 1 | | | | | | | | | | 1 | | | |
| Saltillo | D | 2 | 3 | | | | | | | | | | | | | | | |
| San Luis Potosi | D | 5 | 1 | 9 | 8 | | | | | | | | | | | | | |
| Torreon | D | | | | | | | | | 1 | | 1 | 1 | | | | | |
| Vera Cruz | C | | | | | | | | | | | | | 1 | | 1 | | |
| | D | | | | | | | | | | | | | 1 | | 1 | | |
| Morocco. (See table below.) | | | | | | | | | | | | | | | | | | |
| Nigeria | C | | | 675 | 361 | | | 37 | | 76 | | 74 | | | | | | |
| Nyasaland. (See table below.) | | | | | | | | | | | | | | | | | | |
| Palestine | C | | | 13 | | | | | | | | | | | | | | |
| Paraguay: Asuncion | D | | | 3 | | | | | | | | | | | | | | |
| Persia | C | 89 | 31 | 26 | 20 | | | | 5 | 5 | | 8 | 12 | 25 | | | | |
| | D | 62 | 62 | 9 | 7 | | | | 6 | 1 | | 2 | 1 | 2 | 2 | | | |
| Teheran | C | | | 21 | 17 | 2 | | | 1 | | 3 | 2 | | 6 | | | | |
| | D | | | 8 | 5 | | | | 1 | | 1 | | 1 | 2 | 2 | | | |

| Place | | May 1933 | June 1933 | | | July 1933 | | | August 1933 | | | September 1933 | | | October 1933 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | 1-10 | 11-20 | 21-30 | 1-10 | 11-20 | 21-31 | 1-10 | 11-20 | 21-31 | 1-10 | 11-20 | 21-30 | 1-10 | 11-20 |
| Peru. (See table below.) | | | | | | | | | | | | | | | | |
| Poland | C | 2 | 1 | 1 | | | | | | | | 1 | | 1 | | |
| Portugal (see also table below): | | | | | | | | | | | | | | | | |
| Lisbon | C | 2 | 3 | 5 | 3 | | 2 | 1 | | | | | | | 1 | |
| Oporto | C | 2 | | 3 | 1 | | | | | | 2 | | 1 | 1 | | |
| Siam | D | | | 5 | 1 | | | | | | | | | | | |
| Sierra Leone | C | 51 | 6 | 45 | 6 | [1] 200 | [1] 50 | | | [1] 421 | | [1] 113 | | | | |
| Spain | C | | | 59 | 29 | 11 | 5 | | | 4 | 7 | | 27 | 13 | | |
| Sudan (Anglo-Egyptian) | C | 5 | 1 | 5 | | | 1 | | | | | | | | 14 | 1 |
| Syria: | | | | | | | | | | | | | | | | |
| Beirut | C | 1 | 6 | 14 | 21 | | | | | | | | | | | |
| Provinces | C | 20 | 4 | 9 | 11 | 1 | | 3 | 3 | 16 | | 18 | 2 | 19 | 14 | |
| Turkey. (See table below.) | | | | | | | | | | | | | | | | |
| Union of South Africa: | | | | | | | | | | | | | | | | |
| Cape Province | C | P | | | | | | | | | | | | | | |
| Orange Free State | C | | P | | | | | | | | | | | | | |
| On vessels: | | | | | | | | | | | | | | | | |
| S.S. Rajputana at Aden | C | 1 | | | | | | | | | | | | | | |
| | D | 1 | | | | | | | | | | | | | | |
| S.S. Baron Inchcape at Hong Kong | C | P | | | | | | | | | | | | | | |
| S.S. Fernmoor at Vancouver | C | | 4 | | | | | | | | | | | | | |
| S.S. Egra at Rangoon from Calcutta | C | 1 | | | | | | | | | | | | | | |
| S.S. Arracan at Newport | C | | | | 1 | | | | | | | | | | | |
| S.S. Clan Macquarrie at Suez | C | | | | | 1 | | | | | | | | | | |
| S.S. Sikh at Madras | C | | | | 1 | | | | | | | | | | | |
| S.S. Lichtenfels at Suez from Calcutta | C | | | | | | | | 1 | | | | | | | |
| S.S. Shahjehan at Madras | C | | | | | | 1 | | | | | | | | | |
| S.S. Rohna at Penang from Madras | C | | | | | | | | | | | | 1 | | | |
| Dahomey | C | 11 | 1 | 1 | | 2 | 1 | | 6 | 1 | 1 | 20 | 6 | 1 | 2 | 8 |
| | D | 2 | | 1 | | 1 | | | | 1 | 1 | 5 | 1 | | | 6 |
| Indo-China (see also table above) | C | 149 | 60 | 83 | 54 | 74 | | 31 | 20 | 38 | 21 | | 44 | 37 | 39 | 20 |
| | D | 45 | 16 | 36 | 24 | 31 | | 6 | 5 | 15 | 3 | | 19 | 13 | 15 | 5 |

[1] For 2 weeks.

[2] Dec. 18, 1933: 90 cases of smallpox were reported in Juarez, Mexico, with 18 deaths occurring from Dec. 1 to 16, 1933.

## CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

### SMALLPOX—Continued

[C indicates cases; D, deaths; P, present]

| Place | | May 1933 | June 1933 | July 1933 | August 1933 | September 1933 | October 1933 |
|---|---|---|---|---|---|---|---|
| Bolivia | C | 28 | 12 | 10 | 12 | 15 | 34 |
| Chosen | C | 34 | | 3 | | | |
| | D | 14 | | | | | |
| Costa Rica | C | | | | 2 | | |
| Ecuador | C | [3] 1 | 23 | 12 | | | |
| France | C | 21 | | | | | |
| Greece | C | | | | 1 | | |
| Guatemala | C | 1 | | | | | |

| Place | | May 1933 | June 1933 | July 1933 | August 1933 | September 1933 | October 1933 |
|---|---|---|---|---|---|---|---|
| Mexico (see also table above) | C | | | 9 | 31 | 18 | |
| Morocco | C | 14 | 10 | 8 | 1 | 2 | 3 |
| Nyasaland | C | 303 | 248 | 336 | 347 | 318 | |
| Peru | C | 6 | | 84 | 56 | | |
| Portugal (see also table above) | C | | | | 89 | 105 | |
| | D | | | | 16 | 27 | |
| Turkey | C | | 13 | | 8 | 56 | |

[3] Imported.

### TYPHUS FEVER

[C indicates cases; D, deaths; P, present]

| Place | | Apr. 30–May 27, 1933 | May 28–June 24, 1933 | June 25–July 29, 1933 | Week ended— August 1933 5 | 12 | 19 | 26 | September 1933 2 | 9 | 16 | 23 | 30 | October 1933 7 | 14 | 21 | 28 | November 1933 4 | 11 | 18 |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Algeria: | | | | | | | | | | | | | | | | | | | | |
| Algiers Department | C | • 14 | 11 | 7 | 1 | 1 | | | | | | | | | 1 | | | | | |
| Constantine Department | C | 120 | 157 | 66 | 3 | 8 | | 10 | 2 | 4 | 5 | 1 | 4 | 1 | | 4 | | | | 1 |
| Bone | C | 2 | | 1 | | 1 | | | | | | | | | | | | 1 | | |
| Oran Department | C | | 1 | 2 | | | | | | 2 | | | | | | | 1 | | | |
| Basutoland. (See table below.) | | | | | | | | | | | | | | | | | | | | |
| Bolivia. (See table below.) | | | | | | | | | | | | | | | | | | | | |
| British East Africa: Uganda | C | 10 | 7 | 4 | | 2 | | 2 | 3 | 1 | 5 | 2 | | | 1 | 1 | | | | |
| Bulgaria | C | 8 | 4 | 2 | | 2 | 1 | | | | | 1 | | | | 3 | | 1 | | |
| Chile | C | | | | | | 596 | 432 | 408 | 2,180 | | | | | | | | | | |
| Antofagasta | C | | 1 | | | | | | | | | | | | | | | | | |
| Santiago | C | 134 | 290 | 1,112 | 392 | | | | 124 | 282 | 233 | 218 | 214 | 224 | 292 | 343 | | | | |
| Valparaiso | C | | 1 | 7 | | | 8 | 5 | 5 | 8 | 11 | | 10 | 4 | 8 | 22 | 9 | 5 | 12 | 24 |
| China: | | | | | | | | | | | | | | | | | | | | |
| Hangchow | C | | | | 5 | | | | | | | | | | | | | | | |
| Hankow | C | | | | | | | | | | | | 1 | | | | | | | |
| Harbin | C | | | | | | 8 | 3 | | | | | | | | | | | | |

| | | | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| Nanking | C | | | 1 | | 1 | | | | | | | | 1 | | | | | |
| Shanghai | C | | | 1 | | | 1 | | | | 1 | | | | | | 1 | 1 | |
| Tientsin | C | | | 1 | | | | | | | | | | | | | | | |
| Chosen. (See table below.) | | | | | | | | | | | | | | | | | | | |
| Czechoslovakia. (See table below.) | | | | | | | | | | | | | | | | | | | |
| Egypt: | | | | | | | | | | | | | | | | | | | |
| Alexandria | C | 24 | 17 | 18 | 1 | | | | 1 | | | | | | | | | | |
| Beheira | C | 560 | 569 | 257 | 16 | 21 | 14 | 10 | 7 | 3 | 2 | 4 | 2 | 8 | 5 | 2 | 9 | 9 | 10 |
| Cairo | C | 53 | 28 | 9 | 2 | 1 | | 1 | | | | 1 | | | | | | | |
| Dakahliya | C | 267 | 140 | 82 | 8 | 10 | 19 | 5 | 7 | 2 | 1 | 1 | 3 | | | | | 2 | 2 |
| Damietta | C | | 3 | 3 | | | | 1 | 1 | | | | | | | | | | |
| Gharbiya | C | 448 | 591 | 310 | 22 | 10 | 19 | 28 | 11 | 10 | 8 | 9 | 3 | 1 | 1 | 8 | | 3 | 5 |
| Minufiya | C | | | 52 | 4 | 20 | 4 | 8 | 1 | 1 | 2 | | 2 | 3 | 5 | 2 | 2 | 2 | 3 |
| Qena | C | | | | | | | | 1 | | | 4 | 4 | | | 3 | | 3 | 2 |
| Provinces | C | 1,611 | 1,591 | 785 | 54 | 63 | 56 | 53 | 29 | 22 | 17 | 25 | 14 | 13 | 12 | 15 | 12 | 14 | 29 | 20 |
| Greece. (See table below.) | | | | | | | | | | | | | | | | | | | |
| Guatemala. (See table below.) | | | | | | | | | | | | | | | | | | | |
| Hungary | C | 15 | 11 | | | | | | | | | | | | | 5 | | 10 | | |
| Iraq: Baghdad | C | 2 | 2 | 2 | | 1 | | | | | | | | | | | | | |
| Irish Free State: | | | | | | | | | | | | | | | | | | | |
| Kerry County—Dingle | C | | | | | | | | | | | | | | 4 | | | 4 | |
| Waterford County—Lismore | C | | | | | 1 | | | | | | | | | | | | | |
| Latvia. (See table below.) | | | | | | | | | | | | | | | | | | | |
| Lithuania | C | 31 | 15 | 1 | | 8 | 2 | 1 | | | | | | | | | | | |
| Mexico (see also table below): | | | | | | | | | | | | | | | | | | | |
| Mexico, D.F | C | 15 | 9 | 6 | | 1 | | | 2 | 2 | 2 | | 10 | 5 | 3 | 6 | | 14 | 15 | 10 |
| San Luis Potosí | D | | | | | | | | | | | | 1 | 1 | | | | | | |
| Torreon | C | | | | | | 1 | | | 1 | 1 | | | | | | | 1 | | |
| Morocco | C | 31 | 63 | 14 | 1 | 11 | | | 1 | 1 | | | | 1 | 1 | | | | | |
| Palestine | C | | 1 | 20 | | 2 | | 1 | 1 | 1 | | | 1 | | | | 4 | 4 | | |
| Persia | C | 39 | 25 | 21 | 9 | 4 | 2 | | 4 | 1 | 3 | 6 | 10 | 8 | 5 | 6 | 1 | 5 | | |
| Peru. (See table below.) | | | | | | | | | | | | | | | | | | | | |
| Poland | C | 378 | 322 | 196 | 27 | 19 | 3 | 7 | 6 | 8 | 12 | 8 | 17 | 17 | 8 | 23 | 18 | 34 | 41 | 21 |
| | D | 21 | 13 | 11 | 4 | | 1 | 1 | | | | 1 | | 2 | 1 | | 2 | 3 | | 2 |
| Rumania. (See table below.) | | | | | | | | | | | | | | | | | | | | |
| Spain: Madrid | C | | | | | | | | | 1 | | 1 | | 1 | | | | | | |
| | D | | | | | | | | | | 1 | | | | | | | | | |
| Syria | C | 24 | 21 | 1 | | 1 | | 1 | | | | | | | | | | | | |
| Trans-Jordan | C | 17 | 4 | 2 | | | | | | | | | | | | | | | | |
| Tunisia: | | | | | | | | | | | | | | | | | | | | |
| Tunis | C | | | | | | | | | | 1 | | | | 2 | | | | | |
| Provinces | C | 42 | 65 | 36 | | | | | | | | | | | | | | | | |
| Turkey. (See table below.) | | | | | | | | | | | | | | | | | | | | |
| Union of Socialist Soviet Republics. (See table below.) | | | | | | | | | | | | | | | | | | | | |
| Union of South Africa. (See table below.) | | | | | | | | | | | | | | | | | | | | |
| Yugoslavia. (See table below.) | | | | | | | | | | | | | | | | | | | | |
| On vessels: | | | | | | | | | | | | | | | | | | | | |
| S.S. Chiloe at Antofagasta | C | | 1 | | | | | | | | | | | | | | | | | |
| S.S. Conte Verde at Bombay from Singapore | C | 1 | | | | | | | | | | | | | | | | | | |

## CHOLERA, PLAGUE, SMALLPOX, TYPHUS FEVER, AND YELLOW FEVER—Continued

### TYPHUS FEVER—Continued

[C indicates cases; D, deaths; P, present]

| Place | | May 1933 | June 1933 | July 1933 | August 1933 | September 1933 | October 1933 |
|---|---|---|---|---|---|---|---|
| Basutoland | C | 233 | ----- | 208 | 279 | 259 | ----- |
| Bolivia | C | 98 | 71 | 36 | 34 | 99 | 129 |
| Chosen | C | ----- | ----- | 59 | ----- | ----- | ----- |
| Czechoslovakia | C | 31 | 9 | 6 | 1 | ----- | ----- |
| Greece | C | 12 | 6 | 5 | 4 | 4 | ----- |
| Guatemala | C | 8 | 3 | 1 | 4 | 5 | ----- |
| Latvia | C | ----- | ----- | ----- | 1 | ----- | ----- |
| Mexico (see also table above) | C | ----- | ----- | 12 | 19 | 42 | ----- |
| Peru | C | 28 | 23 | 20 | 60 | ----- | ----- |
| Rumania | C | ----- | ----- | 57 | 22 | 13 | 21 |
| Turkey | C | 16 | 13 | 5 | 7 | 9 | 20 |
| Union of Socialist Soviet Republics | C | 2,051 | ----- | ----- | ----- | ----- | ----- |
| Union of South Africa: | | | | | | | |
| Cape Province | C | 189 | 124 | 195 | 135 | 141 | ----- |
| Natal | C | 6 | 122 | 23 | 18 | 43 | ----- |
| Orange Free State | C | 15 | 34 | 61 | 140 | 180 | ----- |
| Transvaal | C | 1 | 7 | 5 | 3 | 103 | ----- |
| Yugoslavia | C | ----- | 136 | 89 | 33 | 5 | 38 |

### YELLOW FEVER

[C indicates cases; D, deaths; P, present]

| Place | | Apr. 30–May 27, 1933 | May 28–June 24, 1933 | June 25–July 29, 1933 | Week ended— August 1933 | | | | September 1933 | | | | | October 1933 | | | | November 1933 | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| | | | | | 5 | 12 | 19 | 26 | 2 | 9 | 16 | 23 | 30 | 7 | 14 | 21 | 28 | 4 | 11 |
| Brazil: | | | | | | | | | | | | | | | | | | | |
| Ceara State: | | | | | | | | | | | | | | | | | | | |
| Lavras | C | ----- | 1 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- |
| | D | ----- | 1 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- |
| St. Matheus | C | ----- | 1 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- |
| | D | ----- | 1 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- |
| Pernambuco State: | | | | | | | | | | | | | | | | | | | |
| Granito | C | ----- | ----- | 2 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- |
| | D | ----- | ----- | 2 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- |
| Novo Exu [1] | C | ----- | ----- | 1 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- |
| | D | ----- | ----- | 1 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- |
| Salgueiro | C | 1 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- |
| | D | 1 | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- | ----- |

| | | | | | | | | | | | | | | | | | | | |
|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|---|
| French West Africa: | | | | | | | | | | | | | | | | | | | |
| Guinea | C | | | | | | | | | | [2]1 | | | | | | | | |
| Niger Territory | C | | | | | | | 2 | 1 | | | | 2 | 1 | | | | | |
| | D | | | | | | | 2 | | | | | 2 | [3]2 | | | | | |
| Gold Coast | C | 2 | 1 | | | | | | | [2]1 | | | | | [2]1 | | | | 1 |
| | D | | | | | | | | | [2]1 | | | | | | | | | 1 |
| Ivory Coast: | | | | | | | | | | | | | | | | | | | |
| Bouafle | C | | 1 | | | | | | | | | | | | | | | | |
| Gagnoa | C | | [2]1 | | | | | | | | | | | | | | | | |
| | D | | 1 | | | | | | | | | | | | | | | | |
| Nigeria: Kano | C | | | | | | | | | | | | | | | | | 1 | |
| | D | | | | | | | | | | | | | | | | | 1 | |
| Senegal: | | | | | | | | | | | | | | | | | | | |
| Bakel | C | | | | | | | | | | | 2 | | | 1 | | | | |
| | D | | | | | | | | | | | | | | 1 | | | | |
| Kaffrine | C | | | | | | | | | | | | | | 1 | | | | |
| | D | | | | | | | | | | | | | | 1 | | | | |
| St. Louis | D | | | [4]1 | | | | | | | | | | | | | | | |

[1] 2 cases of yellow fever with 2 deaths were reported in Novo Exu, Pernambuco State, Brazil, during the month of June 1933.
[2] Suspected.
[3] Includes 1 suspected death.
[4] Imported.

×

# INDEX

## A


Page

Alabama:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Andervont, H. B.: The specificity of immunity elicited by mouse sarcoma 180 ... 1472
Annual report, abstract of, of the United States marine hospital, Baltimore, Md., for the fiscal year 1933 ... 1496
Anthrax. (*See* Summaries—Disease cases reported monthly by States.)
Argentina: Buenos Aires—Typhus fever—Correction ... 1186
Arizona:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Arkansas:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Armstrong, Charles: Encephalitis, studies on experimental transmission of ... 1341
Arsenoxide content, relation of, to toxicity of fresh and old samples of arsphenamine—Rosenthal and Probey ... 969
Arsphenamine, relation of arsenoxide content to toxicity of fresh and old samples of—Rosenthal and Probey ... 969
Australia: Vital statistics—1932 ... 1414


## B


Bacteria, comparison of the enumeration of, by means of solid and liquid media—Butterfield ... 1292
Badger, L. F.: Rocky Mountain spotted fever, susceptibility of the dog and sheep to the virus of ... 791
Basophilic cells (reticulocytes), estimation of, in blood by examination of ordinary blood film—Jones ... 1011
Bengtson, Ida A.: Seasonal acute conjunctivitis occurring in the Southern States ... 917
Biologic products (establishments licensed for the propagation and sale of viruses, serums, toxins, and analogous products) ... 1401
Blood films for malaria diagnosis, additional notes on the preparation and examination of thick—Komp ... 875
Bloomfield, J. J.:
Preliminary surveys of the industrial environment ... 1343
The size frequency of industrial dusts ... 961
Bone marrow in tularaemia—Lillie and Francis ... 1127
British West Indies: Barbados: Poliomyelitis ... 894
Britten, Rollo H.: Physical impairment and weight (A study of medical examination records of 3,037 men under or over weight for height and age) ... 926
Brundage, Dean K.:
Comparability of sickness records of public utilities ... 1095
Illness among male industrial employees in 1932 as compared with earlier years ... 869
Sickness among male industrial employees during the first quarter of 1933 ... 789
Sickness among male industrial employees during the second quarter of 1933 ... 1191
Butterfield, C. T.:
A note on the food habits of colpidium ... 814
Comparison of the enumeration of bacteria by means of solid and liquid media ... 1292

C

Page

Calcium, effect of, on transplanted mouse tumors—Shear ..... 1103
California:
(*See also* Name of disease—Current weekly State reports.)
(*See also* Summaries—Disease cases reported monthly by States.)
San Benito County—Plague-infected ground squirrels ..... 1050, 1088
Whittier—Plague (human) ..... 1050
Canada:
Ontario: Communicable diseases—Four weeks ended—
June 24, 1933 ..... 894
July 29, 1933 (5 weeks) ..... 1186
August 26, 1933 ..... 1247
September 30, 1933 (5 weeks) ..... 1361
October 28, 1933 ..... 1463
November 25, 1933 ..... 1569
Provinces—Communicable diseases—Two weeks ended—
June 17, 1933 ..... 835
July 1, 1933 ..... 894
July 15, 1933 ..... 989
July 29, 1933 ..... 1092
August 12, 1933 ..... 1124
August 26, 1933 ..... 1186
September 9, 1933 ..... 1247
September 23, 1933 ..... 1306
October 7, 1933 ..... 1361
October 21, 1933 ..... 1429
November 4, 1933 ..... 1484
November 18, 1933 ..... 1510
December 2, 1933 ..... 1569
Quebec—Communicable diseases—Two weeks ended—
May 6, 1933 (10 weeks) ..... 866
July 15, 1933 (10 weeks) ..... 989
July 29, 1933 ..... 1029
August 12, 1933 ..... 1092
August 26, 1933 ..... 1151
September 9, 1933 ..... 1205
September 23, 1933 ..... 1273
October 7, 1933 ..... 1327
October 21, 1933 ..... 1386
November 4, 1933 ..... 1429
November 18, 1933 ..... 1484
December 2, 1933 ..... 1539
Vital statistics—
Year 1932 ..... 1124
First quarter 1933 ..... 1247
Chicken pox. (*See* Summaries—Disease cases reported monthly by States.)
Children, variations of growth in weight of elementary school, 1921–28—Palmer ..... 993
Chile:
Santiago—Typhus fever ..... 990, 1092
Typhus fever ..... 1029
China: Manchuria—Plague ..... 1539
Cholera: Foreign reports ..... 808, 836, 867, 896, 952, 991, 1030, 1056, 1094, 1125, 1152, 1188, 1206, 1250, 1275, 1307, 1328, 1362, 1387, 1414, 1431, 1463, 1486, 1511, 1540, 1571.
Collins, Selwyn D.:
Incidence and clinical symptoms of minor respiratory attacks with special reference to variation with age, sex, and season ..... 1155
Sickness and the economic depression (preliminary report on illness in families of wage earners in Birmingham, Detroit, and Pittsburgh) .. 1251
Colorado:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Colpidium, a note on the food habits of—Butterfield ..... 814

Communicable diseases:
Canada:
Ontario—Four weeks ended— Page
June 24, 1933 894
July 29, 1933 (5 weeks) 1186
August 26, 1933 1247
September 30, 1933 (5 weeks) 1361
October 28, 1933 1463
November 25, 1933 1569
Provinces—Two weeks ended—
June 17, 1933 835
July 1, 1933 894
July 15, 1933 989
July 29, 1933 1092
August 12, 1933 1124
August 26, 1933 1186
September 9, 1933 1247
September 23, 1933 1306
October 7, 1933 1361
October 21, 1933 1429
November 4, 1933 1484
November 18, 1933 1510
December 2, 1933 1569
Quebec—Two weeks ended—
May 6, 1933 (10 weeks) 866
July 15, 1933 (10 weeks) 989
July 29, 1933 1029
August 12, 1933 1092
August 26, 1933 1151
September 9, 1933 1205
September 23, 1933 1273
October 7, 1933 1327
October 21, 1933 1386
November 4, 1933 1429
November 18, 1933 1484
December 2, 1933 1539
Cuba:
Habana—Four weeks ended—
June 17, 1933 835
July 15, 1933 952
August 12, 1933 1124
September 9, 1933 1249
October 7, 1933 1327
November 4, 1933 1463
Provinces:
May 27, 1933 (4 weeks) 835
June 24, 1933 (4 weeks) 952
July 29, 1933 (5 weeks) 1306
September 2, 1933 (5 weeks) 1327
Czechoslovakia:
April 1933 807
May 1933 1029
June 1933 1151
July 1933 1273
August 1933 1386
September 1933 1511
Denmark:
April–June 1933 1054
July 1933 1307
August 1933 1429
Foreign reports 808, 836, 867, 896, 952, 991, 1030, 1056, 1094, 1125, 1152, 1188, 1206, 1250, 1275, 1307, 1328, 1362, 1387, 1414, 1431, 1463, 1486, 1511, 1540, 1571.
Great Britain—England and Wales—Thirteen weeks ended April 1, 1933 836

Communicable diseases—Continued.
Italy—Four weeks ended— Page
March 5, 1933 866
April 2, 1933 1054
April 30, 1933 1187
May 28, 1933 1485
Jamaica—Four weeks ended—
April 22, 1933 1055
May 20, 1933 1093
June 17, 1933 1125
July 15, 1933 1151
August 12, 1933 1187
September 9, 1933 1249
October 7, 1933 1361
November 4, 1933 1430
December 2, 1933 1570
Mexico—Tampico—July 1933 1152
Panama Canal Zone:
April–June 1933 1152
July–September 1933 1430
United States:
Current prevalence of 787, 907, 1069, 1189, 1309, 1417, 1513
Current weekly State reports 800, 829, 860, 888, 945, 982, 1023, 1047, 1085, 1145, 1180, 1198, 1240, 1266, 1299, 1320, 1354, 1379, 1408, 1423, 1456, 1478, 1504, 1533, 1562.
Summary—Disease cases reported monthly by States 802, 831, 862, 890, 947, 984, 1025, 1049, 1087, 1119, 1147, 1182, 1200, 1242, 1268, 1301, 1322, 1381, 1410, 1425, 1458, 1480, 1506, 1535, 1564.
Yugoslavia:
May 1933 808
June 1933 991
July 1933 1205
August 1933 1327
September 1933 1414
October 1933 1511
Conjunctivitis:
(*See* also Summaries—Disease cases reported monthly by States.)
Seasonal acute, occurring in the Southern States—Bengtson 917
Connecticut:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
County health officers, whole-time, 1933 818
Court decisions relating to public health:
Beauty culture law upheld and construed 1297
Birth certificate, certified copy of, used in evidence 1377
Cemeteries, establishment of, held valid 980
City liable for loss of fish in hatchery through pollution of stream 1476
Clerk of courts who is registrar of vital statistics is entitled to fees 1318
Death record, changes in, on file with State health authorities 1178
Death record, issuance of certified copy of, by local health officer 1178
Expense for public health work, bearing by counties of 1502
Food, statute relating to sale of unwholesome 1194
Ice manufactured outside of city, ordinance prohibiting sale of in city of, held valid 1045
Importation of cattle to safeguard against Bang's disease 826
Interfering with township board of health enjoined 1352
Lead poisoning, action to recover damages for injuries from 1238
Liability of city for nuisance caused by effluent from septic tank 1454
Marked milk bottles, city ordinance regarding use of, invalid 1116
Milk and cream, unpasteurized, sale of, held invalid, where ordinance permitted such sale 1560
Milk control board act 884
Morphine, conviction for unlawful possession of, affirmed 858
Narcotic drug law construed 1264
Nuisance, abatement of public, by local board of health 1020

Page

Court decisions relating to public health—Continued.
Occupational disease act upheld 1143
Rocky Mountain spotted fever, compensation for death from 944
Septic tank, damages for injury to land caused by 1421
Sewerage company not liable for damages to land where previous nuisance existed 1406
Vital statistics act 798
Water closets required where sewer connections were possible 1083
Waters, liability for nuisance resulting from impounding of 1194
Cuba:
Cienfuegos—Typhoid fever 1249
Habana—Communicable diseases—Four weeks ended—
June 17, 1933 835
July 15, 1933 952
August 12, 1933 1124
September 9, 1933 1249
October 7, 1933 1327
November 4, 1933 1463
Provinces—Communicable diseases:
May 27, 1933 (4 weeks) 835
June 24, 1933 (4 weeks) 952
July 29, 1933 (5 weeks) 1306
September 2, 1933 (5 weeks) 1327
Current weekly State reports. (*See* Name of disease—Current weekly State reports.)
Czechoslovakia:
Communicable diseases:
April 1933 807
May 1933 1029
June 1933 1151
July 1933 1273
August 1933 1386
September 1933 1510
Vital statistics 1932 1273

D

Davis, Gordon E.: Further studies on the relationship of the viruses of Rocky Mountain spotted fever and Sao Paulo exanthematic typhus 839, 1006
Deaths:
(*See also* Mortality.)
All causes—City reports for week ended—
June 17, 1933 804
June 24, 1933 832
July 1, 1933 863
July 8, 1933 891
July 15, 1933 948
July 22, 1933 985
July 29, 1933 1026
August 5, 1933 1050
August 12, 1933 1089
August 19, 1933 1121
August 26, 1933 1148
September 2, 1933 1183
September 9, 1933 1203
September 16, 1933 1243
September 23, 1933 1269
September 30, 1933 1302
October 7, 1933 1323
October 14, 1933 1357
October 21, 1933 1382
October 28, 1933 1411
November 4, 1933 1426
November 11, 1933 1460
November 18, 1933 1481
November 25, 1933 1507
December 2, 1933 1536
December 9, 1933 1565

Deaths—Continued.
Large cities—Week ended— Page
June 17, 1933 799
June 24, 1933 828
July 1, 1933 859
July 8, 1933 887
July 15, 1933 944
July 22, 1933 981
July 29, 1933 1022
August 5, 1933 1046
August 12, 1933 1084
August 19, 1933 1116
August 26, 1933 1144
September 2, 1933 1179
September 9, 1933 1197
September 16, 1933 1239
September 23, 1933 1265
September 30, 1933 1298
October 7, 1933 1319
October 14, 1933 1353
October 21, 1933 1378
October 28, 1933 1407
November 4, 1933 1422
November 11, 1933 1455
November 18, 1933 1477
November 25, 1933 1503
December 2, 1933 1532
December 9, 1933 1561
Delaware:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Dengue. (See Summaries—Disease cases reported monthly by States.)
Denmark: Communicable diseases:
April–June 1933 1054
July 1933 1307
August 1933 1429
Dermatitis:
An outbreak of, among workers in a rubber manufacturing plant—Schwartz and Tulipan 809
From chemicals used in removing velvet pile—Schwartz and Tulipan 872
Diphtheria:
(*See also* Summaries—Disease cases reported monthly by States.)
City reports for week ended—
June 17, 1933 804
June 24, 1933 832
July 1, 1933 863
July 8, 1933 891
July 15, 1933 948
July 22, 1933 985
July 29, 1933 1026
August 5, 1933 1050
August 12, 1933 1089
August 19, 1933 1121
August 26, 1933 1148
September 2, 1933 1183
September 9, 1933 1202
September 16, 1933 1243
September 23, 1933 1269
September 30, 1933 1302
October 7, 1933 1323
October 14, 1933 1357
October 21, 1933 1382
October 28, 1933 1411
November 4, 1933 1426
November 11, 1933 1460
November 18, 1933 1481
November 25, 1933 1507
December 2, 1933 1536
December 9, 1933 1565

Page

Diphtheria—Continued.
Current weekly State reports ..... 800, 829, 860, 888, 945, 982, 1023, 1047, 1085, 1117, 1145, 1180, 1198, 1240, 1266, 1299, 1320, 1354, 1379, 1408, 1423, 1456, 1478, 1504, 1533, 1562
Directory—State and Insular health authorities, 1933 ..... 1516
District of Columbia:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Drinker, Cecil K.: Zinc in relation to general and industrial hygiene ..... 955
Dusts, the size frequency of industrial—Bloomfield ..... 961
Dusty trades, health of workers in ..... 1398
Dysentery:
(*See also* Summaries—Disease cases reported monthly by States.)
Amoebic, prompt reports of cases of, requested ..... 1401

E

Egypt: Notifiable diseases—Year, 1932 ..... 990
Elvove, Elias: Estimation of fluorides in waters ..... 1219
Employees:
Illness among male industrial, in 1932 as compared with earlier years—Brundage ..... 869
Sickness among male industrial, during the first quarter of 1933 ..... 789
Encephalitis, studies on experimental transmission of—Muckenfuss, Armstrong, and McCordock ..... 1341
Endocarditis, a form of experimental, produced in rabbits—Rose ..... 1038

F

Faget, G. H.: Malarial fever in narcotic addicts; its possible transmission by the hypodermic syringe ..... 1031
Fairhall, Lawrence T.: Zinc in relation to general and industrial hygiene ..... 955
Florida:
(*See* Name of disease—Current weekly State reports.)
(See Summaries—Disease cases reported monthly by States.)
Fluorides in waters, estimation of—Elvove ..... 1219
Food habits of colpidium, a note on the—Butterfield ..... 814
Foreign reports: Communicable diseases ..... 808, 836, 867, 896, 952, 991, 1030, 1056, 1094, 1125, 1152, 1188, 1206, 1250, 1275, 1307, 1328, 1362, 1387, 1414, 1431, 1463, 1486, 1511, 1540, 1571
Francis, Edward: Bone marrow in tularaemia ..... 1127
Fumigants, poisonous, permanent identifying marks on containers of ..... 1419

G

Gas poisoning, methylene blue in the treatment of—Trautman ..... 1443
Georgia:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
German measles. (*See* Summaries—Disease cases reported monthly by States.)
Germany, note on the incidence of endemic goiter in northeastern—Olesen ..... 1074
Goiter, endemic, note on the incidence of, in northeastern Germany—Olesen ..... 1074
Gover, Mary: Incidence and clinical symptoms of minor respiratory attacks with special reference to variation with age, sex, and season ..... 1155
Great Britain:
England and Wales:
Infectious diseases—Thirteen weeks ended April 1, 1933 ..... 836
Vital statistics:
Year 1932 ..... 895
January–March 1933 ..... 836
April–June 1933 ..... 1274
Scotland:
Vital statistics—Quarter ended—
June 30, 1933 ..... 1125
September 30, 1933 ..... 1540
Growth and the economic depression. (A study of the weight of elementary school children in 1921–27 and 1933)—Palmer ..... 1277

H

Page
Health officers:
State and insular, directory, 1933 1516
Whole-time county, 1933 818
Health of workers in a textile plant 1351
Health of workers in dusty trades 1398
Health service, rural, in the United States, extent of, January 1, 1929–December 31, 1932 1222
Heating effect of very high frequency condenser fields on organic fluids and tissues—Schereschewsky 844
Hedley, O. F.: A form of experimental endocarditis produced in rabbits 1038
Himmelsbach, C. K.: Malaria in narcotic addicts at the United States penitentiary annex, Fort Leavenworth, Kans 1465
Hookworm disease. (*See* Summaries—Disease cases reported monthly by States.)

I

Idaho:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Illinois:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Immigrant, the intelligence of the prospective 1115
Immunity elicited by mouse sarcoma 180, the specificity of—Andervont 1472
Impairment, physical, and weight. (A study of medical examination records of 3,037 men under or over weight for height and age)—Britten 926
Indiana:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Industrial employees, male, sickness among—Brundage 789, 869, 1191
Industrial environment, preliminary surveys of the—Bloomfield 1343
Infantile paralysis. (*See* Poliomyelitis.)
Influenza:
(*See also* Summaries—Disease cases reported monthly by States.)
City reports for week ended—
June 17, 1933 804
June 24, 1933 832
July 1, 1933 863
July 8, 1933 891
July 15, 1933 948
July 22, 1933 985
July 29, 1933 1026
August 5, 1933 1050
August 12, 1933 1089
August 19, 1933 1121
August 26, 1933 1148
September 2, 1933 1183
September 9, 1933 1202
September 16, 1933 1243
September 23, 1933 1269
September 30, 1933 1302
October 7, 1933 1323
October 14, 1933 1357
October 21, 1933 1382
October 28, 1933 1411
November 4, 1933 1426
November 11, 1933 1460
November 18, 1933 1481
November 25, 1933 1507
December 2, 1933 1536
December 9, 1933 1565
Current weekly State reports 800, 829, 860, 888, 945, 982, 1023, 1047, 1085, 1117, 1145, 1180, 1198, 1240, 1266, 1299, 1320, 1354, 1379, 1408, 1423, 1456, 1478, 1504, 1533, 1562.
Nicaragua 990

Iowa:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.) Page
Irish Free State: Vital statistics—First and second quarters 1933 1274
Italy: Communicable diseases—Four weeks ended—
March 5, 1933 866
April 2, 1933 1054
April 30, 1933 1187
May 28, 1933 1485

J

Jamaica:
Communicable diseases—Four weeks ended—
April 22, 1933 1055
May 20, 1933 1093
June 17, 1933 1125
July 15, 1933 1151
August 12, 1933 1187
September 9, 1933 1249
October 7, 1933 1361
November 4, 1933 1430
December 2, 1933 1570
Jellison, William L.: Technique for routine and experimental feeding of certain ixodid ticks on guinea pigs and rabbits 1081
Jones, R. R.: Estimation of basophilic cells (reticulocytes) in blood by examination of ordinary blood film 1011

K

Kansas:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Kentucky. (*See* Name of disease—Current weekly State reports.)
Komp, W. H. W.: Additional notes on the preparation and examination of thick blood films for malaria diagnosis 875

L

Lead poisoning:
(*See also* Summaries—Disease cases reported monthly by States.)
In a storage battery plant 1043
Leprosy. (*See* Summaries—Disease cases reported monthly by States.)
Lethargic encephalitis:
(*See also* Summaries—Disease cases reported monthly by States.)
City reports for week ended—
June 17, 1933 806
June 24, 1933 834
July 1, 1933 865
July 8, 1933 893
July 15, 1933 951
July 22, 1933 988
July 29, 1933 1028
August 5, 1933 1053
August 12, 1933 1091
August 19, 1933 1123
August 26, 1933 1150
September 2, 1933 1185
September 9, 1933 1204
September 16, 1933 1246
September 23, 1933 1272
September 30, 1933 1305
October 7, 1933 1326
October 14, 1933 1360
October 21, 1933 1385
October 28, 1933 1413
November 4, 1933 1428
November 11, 1933 1462

Lethargic encephalitis—Continued.
City reports for week ended—Continued. Page
November 18, 1933 1483
November 25, 1933 1509
December 2, 1933 1538
December 9, 1933 1568
In the United States 1120
St. Louis, Mo 1050, 1071, 1088, 1183, 1202, 1243, 1269, 1382
Lillie, R. D.: Bone marrow in tularaemia 1127
Lorenz, Egon: Investigation of mitogenetic radiation by means of a photoelectric counter tube 1311
Louisiana:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)

Mc

McCordock, H. A.: Encephalitis—Studies on experimental transmission 1341
McNamee, Paul D.: Experimental studies of natural purification of polluted waters. VIII. Dissolved oxygen in the presence of organic matter, hypochlorites, and sulphite wastes 1363

M

Maine:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Malaria:
(*See also* Summaries—Disease cases reported monthly by States.)
Diagnosis, additional notes on the preparation and examination of thick blood films for—Komp 875
In narcotic addicts at the United States penitentiary annex, Fort Leavenworth, Kans.—Himmelsbach 1465
Mexico—Monterrey 1362
Therapy, the injection of mosquito sporozoites in—Mayne 909
Malarial fever in narcotic addicts; its possible transmission by the hypodermic syringe—Faget 1031
Maryland:
(*See also* Name of disease—Current weekly State reports.)
(*See also* Summaries—Disease cases reported monthly by States.)
Baltimore, abstract of the annual report of the United States marine hospital at, for the fiscal year 1933 1496
Massachusetts:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Mayne, Bruce: The injection of mosquito sporozoites in malaria therapy 909
Measles:
(*See also* Summaries—Disease cases reported monthly by States.)
City reports for week ended—
June 17, 1933 804
June 24, 1933 832
July 1, 1933 863
July 8, 1933 891
July 15, 1933 948
July 22, 1933 985
July 29, 1933 1026
August 5, 1933 1050
August 12, 1933 1089
August 19, 1933 1121
August 26, 1933 1148
September 2, 1933 1183
September 9, 1933 1202
September 16, 1933 1243
September 23, 1933 1269
September 30, 1933 1302
October 7, 1933 1323
October 14, 1933 1357
October 21, 1933 1382

Measles—Continued.
City reports for week ended—Continued. Page
October 28, 1933 1411
November 4, 1933 1426
November 11, 1933 1460
November 18, 1933 1481
November 25, 1933 1507
December 2, 1933 1536
December 9, 1933 1565
Current weekly State reports 800, 829, 860, 888, 945, 982, 1023, 1047, 1085, 1117, 1145, 1180, 1198, 1240, 1266, 1299, 1320, 1354, 1379, 1408, 1423, 1456, 1478, 1504, 1533, 1562.
Meningococcus meningitis:
(*See also* Summaries—Disease cases reported monthly by States.)
City reports for week ended—
June 17, 1933 806
June 24, 1933 834
July 1, 1933 865
July 8, 1933 893
July 15, 1933 951
July 22, 1933 988
July 29, 1933 1028
August 5, 1933 1053
August 12, 1933 1091
August 19, 1933 1123
August 26, 1933 1150
September 2, 1933 1185
September 9, 1933 1204
September 16, 1933 1246
September 23, 1933 1272
September 30, 1933 1305
October 7, 1933 1326
October 14, 1933 1360
October 21, 1933 1385
October 28, 1933 1413
November 4, 1933 1428
November 11, 1933 1462
November 18, 1933 1483
November 25, 1933 1509
December 2, 1933 1538
December 9, 1933 1568
Current weekly State reports 800, 829, 860, 888, 945, 982, 1023, 1047, 1085, 1117, 1145, 1180, 1198, 1240, 1266, 1299, 1320, 1354, 1379, 1408, 1423, 1456, 1478, 1504, 1533, 1562.
Mercurial poisoning, experimental studies on acute—Rosenthal 1543
Methylene blue in the treatment of HCN gas poisoning—Trautman 1443
Mexico:
Monterrey—Malaria 1362
Tampico—Communicable diseases—July 1933 1152
Vera Cruz:
Notifiable diseases:
December 1932–March 1933 895
April–June 1933 1485
Vital statistics—Year 1932 1030
Michigan:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Miners, coal, mortality of 1420
Minnesota:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Mississippi:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)

Missouri:
(*See also* Name of disease—Current weekly State reports.)
(*See also* Summaries—Disease cases reported monthly by States.) Page
St. Louis—Lethargic encephalitis 1050, 1071, 1088, 1183, 1202, 1243, 1269, 1382
Mitogenetic radiation, investigation of, by means of a photoelectric counter tube—Lorenz 1311
Montana:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Mortality:
(*See also* Deaths.)
From certain causes during the first half of 1933 1447
Of coal miners 1420
Mosquito sporozoites in malaria therapy, the injection of—Mayne 909
Mouse sarcoma 180, the specificity of, elicited by—Andervont 1472
Muckenfuss, Ralph S.: Encephalitis—Studies on experimental transmission 1341
Mumps. (*See* Summaries—Disease cases reported monthly by States.)

N

Narcotic addicts:
Malaria in, at the United States penitentiary annex, Fort Leavenworth, Kans.—Himmelsbach 1465
Malarial fever in: its possible transmission by the hypodermic syringe—Faget 1031
Nebraska:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Neuropsychiatric service in a marine hospital. (Review of 1 year's work of the clinic at Ellis Island)—Reichard 1136
Nevada. (*See* Summaries—Disease cases reported monthly by States.)
New Hampshire:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
New Jersey:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
New Mexico:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
New York:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Nicaragua: Influenza 990
North Carolina:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
North Dakota:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Notifiable diseases:
(*See also* Communicable diseases.)
Egypt—Year 1932 990
Mexico—Vera Cruz:
December 1932–March 1933 895
April–June 1933 1485
Puerto Rico—Four weeks ended—
May 20, 1933 807
June 17, 1933 836
July 15, 1933 990
August 12, 1933 1093
September 9, 1933 1205
September 23, 1933 1275
October 7, 1933 1386
November 4, 1933 1430
December 2, 1933 1540

Notifiable diseases—Continued.
Virgin Islands: Page
February–April 1933 807
May–July 1933 1093
August–October 1933 1486

O

Ohio:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Oklahoma:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Olesen, Robert: Note on the incidence of endemic goiter in northeastern Germany 1074
Oregon:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Organic fluids and tissues, heating effect of very high frequency condenser fields on—Schereschewsky 844

P

Palmer, Carroll E.:
Growth and the economic depression. (A study of the weight of elementary school children in 1921–27 and 1933) 1277
Variations of growth in weight of elementary school children, 1921–28 993
Panama Canal Zone: Communicable diseases:
April–June 1933 1152
July–September 1933 1430
Paratyphoid fever. (*See* Summaries—Disease cases reported monthly by States.)
Parker, R. R.: Further studies on the relationship of the viruses of Rocky Mountain spotted fever and Sao Paulo exanthematic typhus 839, 1006
Patty, F. A.: Acute response of guinea pigs to vapors of some new commercial organic compounds. VII. Dichloroethyl ether 1389
Pellagra:
(*See also* Summaries—Disease cases reported monthly by States.)
City reports for week ended—
June 17, 1933 806
June 24, 1933 834
July 1, 1933 865
July 8, 1933 893
July 15, 1933 951
July 22, 1933 988
July 29, 1933 1028
August 5, 1933 1053
August 12, 1933 1091
August 19, 1933 1123
August 26, 1933 1150
September 2, 1933 1185
September 9, 1933 1204
September 16, 1933 1246
September 23, 1933 1272
September 30, 1933 1305
October 7, 1933 1326
October 14, 1933 1360
October 21, 1933 1385
October 28, 1933 1413
November 4, 1933 1428
November 11, 1933 1462
November 18, 1933 1483
November 25, 1933 1509
December 2, 1933 1538
December 9, 1933 1568
Pennsylvania:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)

Page

Perrott, G. St. J.: Sickness and the economic depression (Preliminary report on illness in families of wage earners in Birmingham, Detroit, and Pittsburgh) 1251
Phenols, tissue, the estimation of—Smith 1487
Philip, C. B.: Technique for routine and experimental feeding of certain ixodid ticks on guinea pigs and rabbits 1081
Photoelectric counter tube, investigation of mitogenetic radiation by means of—Lorenz 1311
Plague (human):
California—Whittier 1050
China—Manchuria 1539
Foreign reports 808, 837, 867, 898, 952, 991, 1030, 1058, 1094, 1126, 1188, 1208, 1250, 1275, 1307, 1330, 1415, 1433, 1464, 1486, 1512, 1541, 1573.
Plague (rodent): California—San Benito County 1050, 1088
Pneumonia (all forms):
City reports for week ended—
June 17, 1933 804
June 24, 1933 832
July 1, 1933 863
July 8, 1933 891
July 15, 1933 948
July 22, 1933 985
July 29, 1933 1026
August 5, 1933 1050
August 12, 1933 1089
August 19, 1933 1121
August 26, 1933 1148
September 2, 1933 1183
September 9, 1933 1202
September 16, 1933 1243
September 23, 1933 1269
September 30, 1933 1302
October 7, 1933 1323
October 14, 1933 1357
October 21, 1933 1382
October 28, 1933 1411
November 4, 1933 1426
November 11, 1933 1460
November 18, 1933 1481
November 25, 1933 1507
December 2, 1933 1536
December 9, 1933 1565
Poland: Vital statistics 1932 1362
Poliomyelitis:
(*See also* Summaries—Disease cases reported monthly by States.)
British West Indies: Barbados 894
City reports for week ended—
June 17, 1933 806
June 24, 1933 834
July 1, 1933 865
July 8, 1933 893
July 15, 1933 951
July 22, 1933 988
July 29, 1933 1028
August 5, 1933 1053
August 12, 1933 1091
August 19, 1933 1123
August 26, 1933 1150
September 2, 1933 1185
September 9, 1933 1204
September 16, 1933 1246
September 23, 1933 1272
September 30, 1933 1305
October 7, 1933 1326
October 14, 1933 1360
October 21, 1933 1385

Poliomyelitis—Continued.
City reports for weeks ended—Continued. Page
October 28, 1933 1413
November 4, 1933 1428
November 11, 1933 1462
November 18, 1933 1483
November 25, 1933 1509
December 2, 1933 1538
December 9, 1933 1568
Current weekly State reports 801, 830, 861, 889, 946, 983, 1024, 1048, 1086, 1118, 1146, 1181, 1199, 1241, 1267, 1300, 1321, 1355, 1380, 1409, 1424, 1457, 1479, 1505, 1534, 1563.
Polluted waters, experimental studies of natural purification in. VIII. Dissolved oxygen in the presence of organic matter, hypochlorites, and sulphite wastes—Theriault and McNamee 1363
Probey, T. F.: Relation of arsenoxide content to toxicity of fresh and old samples of arsphenamine 969
Publications, Public Health Service: (List of publications issued during the period January–June 1933) 1195
Public Health—Court decisions relating to:
Beauty culture law upheld and construed 1297
Birth certificate, certified copy of, used in evidence 1377
Cemeteries, establishment of, held valid 980
City liable for loss of fish in hatchery through pollution of stream 1476
Clerk of courts who is registrar of vital statistics is entitled to fees 1318
Death record, changes in, on file with State health authorities 1178
Death record, issuance of certified copy of, by local health officer 1178
Expense for public health work, bearing by counties of 1502
Food, statute relating to sale of unwholesome 1194
Ice manufactured outside of city, ordinance prohibiting sale of, in city, held valid 1045
Importation of cattle to safe-guard against Bang's disease 826
Interfering with township board of health 1352
Lead poisoning, action to recover damages for injuries from 1238
Liability of city for nuisance caused by effluent from septic tank 1454
Marked milk bottles, city ordinance regarding use of, invalid 1116
Milk and cream, sale of unpasteurized, held invalid where ordinance permitted such sale 1560
Milk control board act 834
Morphine, conviction for unlawful possession of, affirmed 858
Narcotic drug law construed 1264
Nuisance, abatement of public, by local board of health 1020
Occupational disease act upheld 1143
Rocky Mountain spotted fever, compensation for death from 944
Septic tank, damages for injury to land caused by 1421
Sewerage company not liable for damages to land where previous nuisance existed 1406
Vital statistics act 798
Water closets required where sewer connections were possible 1083
Waters, liability for nuisance resulting from impounding of 1194
Public Health Service publications: (List of publications issued during the period January–June 1933) 1195
Public utilities, comparability of sickness records of—Brundage 1095
Puerperal fever. (*See* Summaries—Disease cases reported monthly by States.)
Puerto Rico: Notifiable diseases—Four weeks ended—
May 20, 1933 807
June 17, 1933 836
July 15, 1933 990
August 12, 1933 1093
September 9, 1933 1205
September 23, 1933 1275
October 7, 1933 1386
November 4, 1933 1430
December 2, 1933 1540

## R

Page

Rabies in animals. (*See* Summaries—Disease cases reported monthly by States.)
Rabies in man. (*See* Summaries—Disease cases reported monthly by States.)
Reichard, J. D.: A neuropsychiatric service in a marine hospital (Review of one year's work of the clinic at Ellis Island) ---- 1136
Respiratory attacks, incidence and clinical symptoms of, with special reference to variation with age, sex, and season—Collins and Gover ---- 1155
Rhode Island:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Ringworm. (*See* Summaries—Disease cases reported monthly by States.)
Rocky Mountain spotted (or tick) fever: (*See also* Summaries—Disease cases reported monthly by States.)
Rocky Mountain spotted fever and Sao Paulo exanthematic typhus, further studies on the relationship of the viruses of—Parker and Davis ---- 839, 1006
Rocky Mountain spotted fever: Susceptibility of the dog and sheep to the virus—Badger ---- 791
Rose, Edythe J.: A form of experimental endocarditis produced in rabbits ---- 1038
Rosenthal, Sanford M.:
Experimental studies on acute mercurial poisoning ---- 1543
Relation of arsenoxide content to toxicity of fresh and old samples of arsphenamine ---- 969
Rural health service, extent of, in the United States, January 1, 1929–December 31, 1932 ---- 1222

## S

Scabies. (*See* Summaries—Disease cases reported monthly by States.)
Scarlet fever:
(*See also* Summaries—Disease cases reported monthly by States.)
City reports for week ended—
June 17, 1933 ---- 804
June 24, 1933 ---- 832
July 1, 1933 ---- 863
July 8, 1933 ---- 891
July 15, 1933 ---- 948
July 22, 1933 ---- 985
July 29, 1933 ---- 1026
August 5, 1933 ---- 1050
August 12, 1933 ---- 1089
August 19, 1933 ---- 1121
August 26, 1933 ---- 1148
September 2, 1933 ---- 1183
September 9, 1933 ---- 1202
September 16, 1933 ---- 1243
September 23, 1933 ---- 1269
September 30, 1933 ---- 1302
October 7, 1933 ---- 1323
October 14, 1933 ---- 1357
October 21, 1933 ---- 1382
October 28, 1933 ---- 1411
November 4, 1933 ---- 1426
November 11, 1933 ---- 1460
November 18, 1933 ---- 1481
November 25, 1933 ---- 1507
December 2, 1933 ---- 1536
December 9, 1933 ---- 1565
Current weekly State reports ---- 801, 830, 861, 889, 946, 983, 1024, 1048, 1086, 1118, 1146, 1181, 1199, 1241, 1267, 1300, 1321, 1355, 1380, 1409, 1424, 1457, 1479, 1505, 1534, 1563.
Schereschewsky, J. W.: Heating effect of very high frequency condenser fields on organic fluids and tissues ---- 844

Page

Schrenk, H. H.: Acute response of guinea pigs to vapors of some new commercial organic compounds. VII. Dichloroethyl ether ..... 1389
Schwartz, Louis:
An outbreak of dermatitis among workers in a rubber manufacturing plant ..... 809
Dermatitis from chemicals used in removing velvet pile ..... 872
Septic sore throat. (*See* Summaries—Disease cases reported monthly by States.)
Shear, M. J.: The effect of calcium on transplanted mouse tumors ..... 1103
Sickness among male industrial employees during the first quarter of 1933 ..... 789
Sickness among male industrial employees during the second quarter of 1933—Brundage ..... 1191
Sickness and the economic depression. (Preliminary report on illness in families of wage earners in Birmingham, Detroit, and Pittsburgh)—Perrott, Collins, and Sydenstricker ..... 1251
Sickness records of public utilities, comparability of—Brundage ..... 1095
Smallpox:
(*See also* Summaries—Disease cases reported monthly by States.)
City reports for week ended—
June 17, 1933 ..... 804
June 24, 1933 ..... 832
July 1, 1933 ..... 863
July 8, 1933 ..... 891
July 15, 1933 ..... 948
July 22, 1933 ..... 985
July 29, 1933 ..... 1026
August 5, 1933 ..... 1050
August 12, 1933 ..... 1809
August 19, 1933 ..... 1121
August 26, 1933 ..... 1148
September 2, 1933 ..... 1183
September 9, 1933 ..... 1202
September 16, 1933 ..... 1243
September 23, 1933 ..... 1269
September 30, 1933 ..... 1302
October 7, 1933 ..... 1323
October 14, 1933 ..... 1357
October 21, 1933 ..... 1382
October 28, 1933 ..... 1411
November 4, 1933 ..... 1426
November 11, 1933 ..... 1460
November 18, 1933 ..... 1481
November 25, 1933 ..... 1507
December 2, 1933 ..... 1536
December 9, 1933 ..... 1565
Current weekly State reports ..... 801, 830, 861, 889, 946, 983, 1024, 1048, 1086, 1118, 1146, 1181, 1199, 1241, 1267, 1300, 1321, 1355, 1380, 1409, 1424, 1457, 1479, 1505, 1534, 1563.
Foreign reports ..... 900, 991, 1061, 1210, 1332, 1436, 1576
Smith, Maurice I.: The estimation of tissue phenols ..... 1487
South Carolina:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
South Dakota:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
State and insular health authorities, directory of, 1933 ..... 1516
Straits Settlements: Singapore—Vital statistics 1932 ..... 1249
Streams, studies of the pollution and natural purification of. (A resurvey of the Ohio River between Cincinnati, Ohio, and Louisville, Ky.) ..... 976
Summaries: Communicable diseases—Disease cases reported monthly by State health officers ..... 802, 831, 862, 890, 947, 984, 1025, 1049, 1087, 1119, 1147, 1182, 1200, 1242, 1268, 1301, 1322, 1356, 1381, 1410, 1425, 1458, 1480, 1506, 1535, 1564.

Page

Sydenstricker, Edgar: Sickness and the economic depression (Preliminary report on illness in families of wage earners in Birmingham, Detroit, and Pittsburgh) 1251

T

Tennessee:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Tetanus. (*See* Summaries—Disease cases reported monthly by States.)
Texas:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Textile plant, health of workers in a 1351
Theriault, Emery J.: Experimental studies of natural purification in polluted waters. VIII. Dissolved oxygen in the presence of organic matter, hypochlorites and sulphite wastes 1363
Ticks, ixodid, on guinea pigs and rabbits, technique for routine and experimental feeding of certain—Jellison and Philip 1081
Tissue phenols, the estimation of—Smith 1487
Trachoma. (*See* Summaries—Disease cases reported monthly by States.)
Trautman, J. A.: Methylene blue in the treatment of HCN gas poisoning 1443
Trinidad: Port of Spain—Vital statistics, 1932 1511
Tuberculosis:
City reports for week ended—
June 17, 1933 804
June 24, 1933 832
July 1, 1933 863
July 8, 1933 891
July 15, 1933 948
July 22, 1933 985
July 29, 1933 1026
August 5, 1933 1050
August 12, 1933 1089
August 19, 1933 1121
August 26, 1933 1148
September 2, 1933 1183
September 9, 1933 1202
September 16, 1933 1243
September 23, 1933 1269
September 30, 1933 1302
October 7, 1933 1323
October 14, 1933 1357
October 21, 1933 1382
October 28, 1933 1411
November 4, 1933 1426
November 11, 1933 1460
November 18, 1933 1481
November 25, 1933 1507
December 2, 1933 1536
December 9, 1933 1565
Tularaemia:
(*See also* Summaries—Disease cases reported monthly by States.)
Bone marrow in—Lillie and Francis 1127
Tulipan, Louis:
An outbreak of dermatitis among workers in a rubber manufacturing plant 809
Dermatitis from chemicals used in removing velvet pile 872
Tumors, effect of calcium on transplanted mouse—Shear 1103
Typhoid fever:
(*See also* Summaries—Disease cases reported monthly by States.)
City reports for week ended—
June 17, 1933 804
June 24, 1933 832
July 1, 1933 863
July 8, 1933 891
July 15, 1933 948

Typhoid fever—Continued.
City reports for week ended—Continued. Page
July 22, 1933 985
July 29, 1933 1026
August 5, 1933 1050
August 12, 1933 1089
August 19, 1933 1121
August 26, 1933 1148
September 2, 1933 1183
September 9, 1933 1202
September 16, 1933 1243
September 23, 1933 1269
September 30, 1933 1302
October 7, 1933 1323
October 14, 1933 1357
October 21, 1933 1382
October 28, 1933 1411
November 4, 1933 1426
November 11, 1933 1460
November 18, 1933 1481
November 25, 1933 1507
December 2, 1933 1536
December 9, 1933 1565
Cuba—Cienfuegos 1249
Current weekly State reports 801, 830, 861, 889, 946, 983, 1024, 1048, 1086, 1118, 1146, 1181, 1199, 1241, 1267, 1300, 1321, 1355, 1380, 1409, 1424, 1457, 1479, 1505, 1534, 1563.
Typhus fever:
(*See also* City reports.)
Argentina—Buenos Aires—Correction 1186
Chile 1029
Santiago 990, 1092
Foreign reports 837, 867, 904, 953, 1030, 1066, 1126, 1152, 1188, 1215, 1274, 1337, 1362, 1387, 1415, 1440, 1580
(Experimental transmission of endemic typhus by Xenopsylla astia)—Workman 795
Typhus, Sao Paulo exanthematic, and Rocky Mountain spotted fever, further studies on the relationship of the viruses of—Parker and Davis 839, 1006

U

Undulant fever. (*See* Summaries—Disease cases reported monthly by States.)
United States:
Communicable diseases:
Current prevalence of 787, 907, 1069, 1189, 1309, 1417, 1513
Current weekly State reports 800, 829, 860, 888, 945, 982, 1023, 1047, 1085, 1145, 1180, 1198, 1240, 1266, 1299, 1320, 1354, 1379, 1408, 1423, 1456, 1478, 1504, 1533, 1562.
Summaries—Disease cases reported monthly by State health officers 802, 831, 862, 890, 947, 984, 1025, 1049, 1087, 1119, 1147, 1182, 1200, 1242, 1268, 1301, 1322, 1356, 1381, 1410, 1425, 1458, 1480, 1506, 1535, 1564.
Deaths—Large cities—Week ended—
June 17, 1933 799
June 24, 1933 828
July 1, 1933 859
July 8, 1933 887
July 15, 1933 944
July 22, 1933 981
July 29, 1933 1022
August 5, 1933 1046
August 12, 1933 1084
August 19, 1933 1116

United States—Continued.
Deaths—Large cities—Week ended—Continued. Page
August 26, 1933 1144
September 2, 1933 1179
September 9, 1933 1197
September 16, 1933 1239
September 23, 1933 1265
September 30, 1933 1298
October 7, 1933 1319
October 14, 1933 1353
October 21, 1933 1378
October 28, 1933 1407
November 4, 1933 1422
November 11, 1933 1455
November 18, 1933 1477
November 25, 1933 1503
December 2, 1933 1532
December 9, 1933 1561
Lethargic encephalitis in the 1120
Utah: (*See* Name of Disease—Current weekly State reports.)

V

Vapors of some new commercial organic compounds, acute response of guinea pigs to. VII. Dichloroethyl ether—Schrenk and Patty 1389
Venezuela: Vital statistics, year 1932 866
Vermont:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Vincent's angina: (*See* Summaries—Disease cases reported monthly by States.)
Virginia:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Virgin Islands:
Notifiable diseases:
February–April 1933 807
May–July 1933 1093
August–October 1933 1486
Vital statistics:
Australia, 1932 1414
Canada:
Year 1932 1124
First quarter, 1933 1247
Czechoslovakia, 1932 1273
Great Britain:
England and Wales:
Year, 1932 895
January–March, 1933 836
April–June 1933 1274
Scotland—Quarter ended—
June 30, 1933 1125
September 30, 1933 1540
Irish Free State—First and second quarters, 1933 1274
Mexico—Vera Cruz, 1932 1030
Poland, 1932 1362
Straits Settlements—Singapore, 1932 1249
Trinidad: Port of Spain, 1932 1511
Venezuela, 1932 866

W

Washington:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Weight and physical impairment. (A study of medical examination records of 3,037 men under or over weight for height and age.)—Britten 926

West Virginia:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Whooping cough:
(*See also* Summaries—Disease cases reported monthly by States.)
City reports for week ended— Page
June 17, 1933 804
June 24, 1933 832
July 1, 1933 863
July 8, 1933 891
July 15, 1933 948
July 22, 1933 985
July 29, 1933 1026
August 5, 1933 1050
August 12, 1933 1089
August 19, 1933 1121
August 26, 1933 1148
September 2, 1933 1183
September 9, 1933 1202
September 16, 1933 1243
September 23, 1933 1269
September 30, 1933 1302
October 7, 1933 1323
October 14, 1933 1357
October 21, 1933 1382
October 28, 1933 1411
November 4, 1933 1426
November 11, 1933 1460
November 18, 1933 1481
November 25, 1933 1507
December 2, 1933 1536
December 9, 1933 1565
Wisconsin:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)
Workman, W. G.: Typhus fever. (Experimental transmission of endemic typhus by Xenopsylla astia) 795
Wyoming:
(*See* Name of disease—Current weekly State reports.)
(*See* Summaries—Disease cases reported monthly by States.)

## Y

Yellow fever: Foreign reports 808, 837, 906, 1068, 1153, 1217, 1250, 1274, 1339, 1442, 1464, 1486, 1512, 1541, 1582
Yugoslavia: Communicable diseases—
May 1933 808
June 1933 991
July 1933 1205
August 1933 1327
September 1933 1414
October 1933 1511

## Z

Zinc in relation to general and industrial hygiene—Drinker and Fairhall 955

○